

Analysis of Physio-Chemical Properties and Plant Growth Characteristics using Different Growing Media

Abstract

Soilless culture is a technique of crop production using no soil. The main reason towards need for soil to soilless culture for horticultural crops is the problem related to proliferation of soil borne pathogen in the soil cultivation. Recent research studies reported that commercial production of greenhouse vegetables with soilless media has paved way for reduction in economic losses caused by soil-borne pathogens. In this context, the present study was initiated to identify suitable substrate mixes for soilless culture. The experiment was laid out in a randomized block design. There were fifteen different media levels including control, with three replications. Crop Tomato TNAU hybrid CO3 were assessed for the mean performance in respect of growth characters namely shoot length, root length, germination percent and vigour index. The treatment Vermicompost: Coirpith (3:1) was found to have the highest value for growth characters for the crop. In the point of physical and chemical properties, all combination media almost showed optimal range for crop growth. Further a cost analysis was made to find the suitable media based on cost aspects comparing reduction in vigour index and cost of media with that of best media. Treatment T₇ (Vermicompost: coir pith (2:1)) and T₆ (Vermicompost: Coir pith (1:3)) were found to be the best media considering above factors.

Keywords: Physio-Chemical properties, plant growth characteristics, soilless growing media, cost analysis

Introduction

Soil-based agriculture is facing some major challenges with the advent of civilization all over the world, such as decrease in per capita land availability, non-availability of work force, incidence of pests and diseases, etc. Apart from this, due to rapid urbanization and industrialization as well as threats from climate change and its related adverse effects, the land cultivation is going to face further challenging threats. Under such circumstances, in the near future it becomes intricate to feed entire population using the production from soil field system. Naturally, soilless culture is gaining its relevance in the present scenario, to cope-up with these challenges. Soilless culture is the modern cultivation system of plants that use either inert organic or inorganic substrate through nutrient solution nourishment. Possibly it is the most intensive culture system utilizing all the resources efficiently for maximizing yield

of crops and the most intense form of agricultural enterprises for commercial production of vegetables. Several studies suggested soilless culture is an alternative to traditional field production for high-value vegetable crops. Md. Asaduzzaman et al., 2017 reported that the quality of horticultural crops grown through soilless culture improves significantly compared to conventional soil culture. This artificial growing system provides plants with mechanical support, water and mineral nutrient for higher growth and development. Soilless growing media are easier to handle and it may provide better growing environment (in terms of one or more aspects of plant growth) compared to soil culture. Organic substrates includes sawdust, coco peat, vermicompost, peat moss, woodchips, fleece, marc, bark, etc. whereas, inorganic substrate of natural origin are perlite, vermiculite, zeolite, gravel, rock wool, sand, glass wool, pumice, sepiolite, expanded clay, volcanic tuff and synthetically produced substrates are hydrogel, foam mates (polyurethane), oasis (plastic foam), etc. Various raw materials have been used to produce growing media for vegetable production throughout the world. The present research was initiated in this context to analysis the Physio-chemical properties of different growing media, to conduct germination studies to find the best media and to assess the economical viability of growing media to establish suitable substratum mixes for soilless cultivation.

Methodology

The field experiment was conducted in the farm Imayam Institute of Agriculture and Technology, Thuraiyur. The experimental site lies geographically in between 11° 8' 29" N latitude and 78° 35' 40" E longitude and at an altitude of 151 m above mean sea level. The experiment was designed under Randomized block design as 15 treatments with three replications as mentioned in Table (1). The loamy soil available was used as control. Physico chemical prosperities of media characteristics like bulk density, particle density, porosity, pH and EC were analyzed by standard procedure reported by Keen and Rackzowski, 1921. The climatic parameters such as temperature and relative humidity are monitored at 8.30 am and 2.30 pm with the digital thermo hygrometer throughout the crop growth period of 09.03.2019 to 08.04.19.

Table.1. Media and their combinations for different treatments

Treatment	Growing media(v/v)
T ₁	Coir pith
T ₂	Sawdust
T ₃	Vermicompost
T ₄	Vermicompost : Coir pith (1:1)
T ₅	Vermicompost : Coir pith (1:2)
T ₆	Vermicompost : Coir pith (1:3)
T ₇	Vermicompost : Coir pith (2:1)
T ₈	Vermicompost : Coir pith (3:1)

T ₉	Vermicompost : Sawdust (1:1)
T ₁₀	Vermicompost : Sawdust (1:2)
T ₁₁	Vermicompost: Sawdust(1:3)
T ₁₂	Vermicompost : Sawdust (2:1)
T ₁₃	Vermicompost : Sawdust (3:1)
T ₁₄	Vermicompost : Coir pith: Sawdust (1:1:1)
T ₁₅	Control (Clay loam)

Hybrid variety of Tomato crop (CO3) 20 days seedlings with three replications were taken for the study and their physiological parameters and growth characteristics were determined as follows.

Root length

Root length was measured from the collar region to the tip of the primary root and the average expressed in centimeters.

Shoot length

The length from the collar region to the tip of the shoot was measured as shoot length and the average was expressed in centimeters.

Germination percentage (GP)

The germination percentage was calculated based on formula

$$GP = \frac{\text{No. of sprouted seed}}{\text{No. of total seeds}} \times 100 \dots \dots \dots (1)$$

Vigour index (VI)

$$VI = GP \times (\text{Root length, cm} + \text{Shoot length, cm}) \dots \dots \dots (2)$$

Statistical analysis

Statistical analysis (AGRES) was carried out to study the effects of alternate growing systems in plant growth parameters. Least significant differences test (LSD) was used to compare the significant differences among mean of the treatments at 0.05 level of probability.

Cost analysis

Cost analysis was used to find out the best suitable media which has the lowest % reduction in vigour index along with percentage reduction in total cost.

Results and Discussion

The study was aimed to identify a suitable growing media. In this context the Physico-chemical characteristics of growing media were analyzed and the environmental factors like temperature and relative humidity were recorded. The effects of growing media on plant growth character were observed. The cost analyses were done to determine the best media for crop growth. The observations of the study were analyzed and the results are as follows.

Selection of suitable growing media based on physico-chemical properties, climatic parameters, physiological parameters

Bulk density

Bulk density of medium is one of the indices for the evaluation of physical conditions of the growing media. Optimum bulk density for optimal plant growth ranges between 0.25 to 0.58 g cm⁻³ and it is high as 1.3 g cm⁻³ Conover, 1967. In this study the combination of Vermicompost : Coir pith (T₈ - 3:1) was observed with bulk density of 0.55 g cm⁻³ was in the recommended range. The highest value of bulk density was observed for loamy soil (1.3 g cm⁻³) and lowest bulk density was noticed in coir pith (0.28 g cm⁻³) as individual media but in combination, T₉ - Vermicompost : Sawdust (1:1) recorded the lowest value (0.35 g cm⁻³).

Porosity characteristics

The porosity characteristics of a media are measured in terms of Total Porosity (TP), Air Filled Porosity (AFP) and Water Holding Capacity (WHC). The highest porosity (88%) was obtained for coir pith followed by Vermicompost: Coir pith (3:1) (78%). Loamy soil had low porosity 40 %. But all combination media almost showed optimal total porosity (65 – 80 %), air filled porosity (20 - 30%) and water holding capacity (45- 55%). These results are in conformity with the results reported by Di Lorenzo *et al.*, 2013. Raviv and Lieth 2008 reported that most media and mixes have an air filled porosity of (10-30 %). The results obtained in this experiment are similar and all media combinations employed as treatments have shown ideal porosity characteristics.

pH value variation in treatments

The values were all in slightly acidic or neutral range. Except for control none of the media showed pH in the alkaline ranges. The highest pH was observed in treatments, T₃ - (7.32) and in control (7.57) but in the case of media combination the highest pH was observed in T₁₃ (3:1) (7.05), the lowest value of pH was observed in the case of media T₁ - (6.1). The pH is an important factor in the availability and the uptake of nutrients and satisfactory pH of the medium ranged between 5.5 and 6.5 as reported by Waters *et al.*, 1970.

EC value variation in treatments

The EC value of the different media varied between 0.13 to 0.45 dS m⁻¹ during the growing period. The maximum EC value of 0.45 dS m⁻¹ was recorded for T₃ (Vermicompost) and the lowest value for T₂- Sawdust (0.13 dS m⁻¹). According to the results of Bunt 1988 there was no salinity problem in the substrates. **Substrate should have low salinity because it decreases mineral acquisition and salts that enter the soil decrease the capacity of roots to absorb water by decreasing the potential gradient from the soil to the roots.** The EC values were below the established limit (0.5 dS m⁻¹) for an ideal substrate reported by Abad *et al.*,

2001. The low pH and EC values might have been ideal for crop growth whereas the high values recorded for these parameters might have led to reduced plant growth as already reported by Francesco *et al.*, 2005.

Variation of temperature and relative humidity during the growth period

The maximum minimum temperature and relative humidity 38.5°C, 26.0°C and 79 %, 26 % respectively was recorded during the crop growing period. For most of the days, the temperature exceeded 35°C during the recorded time.

Ganesan, 2002 reported that the average 24 hr temperature is believed to be responsible for the growth rate of the crop the higher the average air temperature, the faster the growth. Higher humidity was observed during morning hours and gradually decreased in the afternoon because of increase in temperature. Growth is positively correlated to relative humidity and normal plant growth occurs at relative humidity of 25 - 80 percent.

Effect of different growing media on plant growth parameters

Significant differences were observed in the mean shoot length, root length, germination percentage and vigour index between treatments. Treatment T₈ (Vermicompost: Coir pith- 3:1) and treatment T₂ (Sawdust) recorded the highest and lowest root length, shoot length, germination percentage and vigour index as 18 cm, 9.0cm, 80.11%, 3.18 and 8.1cm, 3.2cm, 36.13%, 2.47 respectively, 30 days after sowing (DAS). The variations are graphically illustrated in Figure (1 to 4).

The significant increase in plant growth in treatment T₈ (Vermicompost: Coirpith - 3:1) may be because of the good physical properties of vermicompost and due to high organic matter, macro and micronutrient content present in it. The T₈ and T₂ treatments were shown in Figure 5 and 6 which was taken on 30 days after sowing. Similar results were obtained by Umamaheswari, 2005 who reported that the vermicompost is the granular aggregate, the stability of which is due to the mucopolysaccharides of microbes and earthworms and it is rich in plant nutrient.

Cost analysis of growing media

Based on all treatments conducted T₈ displayed more positive influence on plant growth but was costly based on economic aspects (Rs. 4370.0). As compared to T₈, treatments T₇ and T₆ shows the lowest reduction in vigour index and highest reduction in cost (2.3 %), (4.3 %) and (8.5%), (45.3%) respectively. Based on the above analysis regarding performance of different media T₆ and T₇ may be suggested as suitable and economically viable media.




Figure 1 Effect of different growing media on shoot length (cm)




Figure 2 Effect of different growing media on Root length (cm)




Figure 3 Effect of different growing media on germination percentage




Figure 4 Effect of different growing media on vigour index



Figure 5 Treatment T₈



Figure 6 Treatment T₂

Table 1. Percentage reduction in vigour index and increase or decrease in cost compared with best media, T₈ (Vermicompost: Coir pith - 3:1)

Treatments	Vigour index (VI)	% Reduction in VI	Total cost /m ³	% Increase or decreases in Total cost
T ₁	2.77	12.9	1400.0	-68.0
T ₂	2.47	22.3	1020.0	-76.7
T ₃	3.02	4.8	5360.0	22.7

T ₄	2.91	8.5	3380.0	-22.7
T ₅	2.80	11.7	2692.8	-38.4
T ₆	3.04	4.3	2390.0	-45.3
T ₇	3.10	2.3	3999.6	-8.5
T ₈	3.18	0.0	4370.0	0.0
T ₉	2.96	6.9	3190.0	-27.0
T ₁₀	2.86	10.1	2442.0	-44.1
T ₁₁	2.98	6.3	2105.0	-51.8
T ₁₂	2.84	10.7	3874.2	-11.3
T ₁₃	2.73	14.0	4275.0	-2.2
T ₁₄	2.77	12.9	2567.4	-41.2
T ₁₅	2.63	17.2		

Conclusion

The plant growth was high in treatments in which coirpith as combination media, because coirpith had good physical properties such as high water holding capacity, good aeration and excellent thermal properties and it enhanced root and shoot development throughout crop period. The treatment T₂ and treatments in combination with sawdust showed poor performance compared to other media. The results of the present study showed that cultivating crops in a growing medium consisting of Vermicompost : Coir pith (3:1) can highly influence the plant growth and increase the yield. Based on cost analysis T₇ (Vermicompost : Coir pith (2:1)) and T₆ (Vermicompost : Coir pith (1:3)) were suggested as suitable and economically viable media because compared with other media and combinations T₇ and T₆ showed less reduction in vigour index (2 - 4%) and less cost (3999.00/m³ and 2390.00/m³). Treatment T₇ (Vermicompost: coir pith (2:1)) and T₆ (Vermicompost: Coir pith (1:3)) were found to be the best media considering above factors.

References

- Abad, M., Noguera, P. and Bures, S. 2001. National inventory of organic wastes for use as growing media for ornamental potted plant production, case study in Spain. **Biores. Technol.**77:197-200.
- Bhardwaj, R. L. 2014. Effect of growing media on germination and seedling growth of papaya cv. „Red lady“. **African journal of plant science**, Vol. 8(4): 174-188
- Bunt, A. C. 1988. Media and Mixes for Container- Grown Plants. 1st Edn., **Springer, London.**
- Conover, C. A. 1967. Soil mixes for ornamental plants. **Florida Flower Grower**, 4: 1-4.
- Di Lorenzo, R., Pisciotta, A., Santamaria, P. and Scariot, V. 2013. From Soil to Soilless in Horticulture: Quality and Typicity. **Italian Journal of Agronomy**, Vol 8(4)

Ganesan, M. 2002. Effect of Poly-Greenhouse Models on Plant Growth and Yield of Tomato (*Lycopersicon esculentum*). **JRD Tata Ecotechnology Centre, M. S. Swaminathan Research Foundation.**

Francesco Montesano., Gianfranco favuzzi., Angelo parente and Francesco serionad Pietro santamaria. 2005. NFT versus subirrigation: I. yield and water use efficiency of tomato grown in closed soilless systems under salinity or water stress. **Hort. Sci., 40(4): 1011-1012.**

Jozef Nurzynski. 2013. Effect of Substrates on Nutrient Content in Root Zone and Leaves of Greenhouse Tomato. **Acta Sci. Pol., Hortorum Cultus 12(5): 169-178.**

Keen, B.A. and Raczkowski, H. (1921). Relation between the clay content and certain physical properties of a soil. **Journal of Agricultural Science, 11: 441-449**

Meena, A.K., O.P. Garhwal, Arun Kumar Mahawar and Singh, S.P. 2017. Effect of Different Growing Media on Seedling Growth Parameters and Economics of Papaya (*Carica papaya* L) cv. *Pusa Delicious*. **Int.J.Curr.Microbiol.App.Sci. 6(6): 2964-2972.**

Md Asaduzzaman, Md. Saifullah, AKM Salim Reza Mollick, Md. Mokter Hosssain, GMA Halim and Toshiki Asao. 2015. Soilless culture – Use of Substrates for the Production of Quality Horticultural Crops. Published by **In Tech. 1(1)**

Raviv, M. and Lieth, J. 2008. Soilless culture: theory and practice, **Elsevier Science Ltd.**

Theodore T. Kozlowski, Stephen G. Pallardy, 1997. Environmental Regulation of Vegetative Growth. **Growth Control in Woody Plants. 4(14)**

Umamaheswari, S. 2005. Vermicomposting of mango litter using earthworm perionyx excavates. **J. Ecotoxicol environmental monitoring. 15(1): 69-72**

Umesha, B., Bosu, S. S., Rema, K.P. and Murumkar, A.R. 2013. Drip irrigation with fertigation in soil-less media for tomato under controlled cultivation. **Journal of Applied Horticulture, 15(3): 195-197**

Yaseer Suhaimi, M., Mohamad, A. M., Mahamud S. and Khadzir, D. 2012. Effects of substrate on growth and yield of ginger cultivated using soilless culture.