

**Anti-oxidant and Anti-inflammatory activity of ethyl acetate fraction of  
*Moringa oleifera* flowers**

**A. Rajeshkanna<sup>1,2\*</sup>, M. M. Senthamilselvi<sup>2,3</sup> D. Prabhakaran<sup>2,4</sup>**

<sup>1</sup>Department of Chemistry, Ananda College, Devakottai, Tamil Nadu, India.

<sup>2</sup>Department of Chemistry, Periyar E.V.R College, Affiliated to Bharathidasan University,  
Trichy, Tamil Nadu, India.

<sup>3</sup>Principal (Rted), Government Arts College, Surandai , Tirunelveli Dt, Tamil Nadu, India.

<sup>4</sup>Chettinad Cement Corporation Ltd., Ariyalur, Tamil Nadu, India.

**\*Corresponding author:**

A.Rajeshkanna,

Asst professor,

Department of Chemistry,

Ananda College, Devakottai, Tamil Nadu, India,

***E.mail id:*** [a.rajeshkannachem@gmail.com](mailto:a.rajeshkannachem@gmail.com)

## **ABSTRACT**

**Aims :** To investigate the antioxidant and anti-inflammatory activities of ethyl acetate extract of *Moringa oleifera* flowers.

**Place and duration of study :** The research work was carried out at Research laboratory, Department of chemistry, Periyar E.V.R College, Trichy-23, between April 2017 and January 2018.

**Methodology:** Extraction and fractionation was carried out from the solvents of ethanol, benzene, petroleum ether, diethyl ether and ethyl acetate. The anti-inflammatory effect of the extract was investigated by HRBC membrane stabilization and Albumin denaturation methods. Anti-oxidant effect of the extract was determined by DPPH assay and ABTS method.

**Results:** The dry sample extracted from the ethyl acetate fraction of *Moringa oleifera* flowers possesses highly anti-oxidant activity showed by the DPPH assay and ABTS method and also having anti-inflammatory activity is determined by human red blood cell (HRBC) membrane stabilization and Albumin denaturation methods. However, these effects need to be confirmed using *in vivo* models and clinical trials before its utilization as a therapeutic agent.

**Conclusion:** The present study was concluded that the dry sample of ethyl acetate fraction of *Moringa oleifera* flowers possesses effective anti-oxidant and anti-inflammatory activities.

**Keywords:** ABTS assay, Anti-inflammatory activity, Antioxidant activity, HRBC method, *Moringa oleifera*.

## **1. INTRODUCTION**

Peoples may be exposed to endogenous free radicals sources like Ultraviolet radiation, cigarette and vehicles smoke, atmospheric pollutants, industrial effluent etc, in our day to day life<sup>[1]</sup>. Free radicals cause several metabolic disorders in cells in the human body. Some of the oxidants like reactive oxygen species reactive nitrogen species, hydroxyl radical and hydrogen peroxide may produce cancer, cardiovascular disease, and cataract, slow down the ageing in human beings. So we protect from these diseases intake of antioxidants from the external sources<sup>[2]</sup>. Antioxidents are the substances that act against oxidants<sup>[3]</sup>. The high potential antioxidants such as vitamin C, vitamin E,  $\beta$ - Carotene, flavonoids and polyphenols are available in natural

plants<sup>[4]</sup>. As oxidative strain is a vital role in liver pathologies and their progression, the use of antioxidants in a natural remedies neutralize the liver damage<sup>[5]</sup>.

Inflammation is a normal body response to a traumatic injury, irritation from harmful chemicals, or infection caused by microbial pathogens. It leading to painful conditions like rheumatoid arthritis, asthma, allergy, inflammatory bowel syndrome and atherosclerosis are caused by the presence of plasma and white blood cells in the affected area<sup>[6]</sup>. Pain and inflammatory responses in the peripheral and central nervous systems play key roles in the growth and persistence of many pathological pain states<sup>[7]</sup>. A variety of natural compounds are able to lightening the inflammation by reducing the formation of inflammatory mediators, or modulating inflammatory<sup>[8]</sup>.

*Moringa oleifera* tree belongs to Moringaceae family<sup>[9]</sup>. It is widely cultivated species in home gardens and sub-Himalayan tracts of India, Pakistan, Bangladesh and Afghanistan<sup>[10]</sup> for the purposes of food and medicines by the presence of vitamins, proteins and phenolic compounds in various parts<sup>[11]</sup>. The previous researches reported all the plant parts were contains different variety of biochemical compounds with anticarcinogenic<sup>[10]</sup>, anti-inflammatory<sup>[12]</sup>, antidiabetic<sup>[13]</sup>, antioxidant<sup>[14]</sup>, and antimicrobial effects<sup>[15]</sup>. However it is known that phytochemical concentration varies from part to part of the plant. To the best of our knowledge there have not been any phytochemical studies focusing on the flowers of *M. oleifera*. Thus this study set out to investigate the anti-oxidant and anti-inflammatory activities of this plant part as a possible clinical therapeutic agent.

## **2. MATERIALS AND METHODS**

### **2.1 Collection of flowers**

Fresh flowers of *M. oleifera* were collected from Karaikudi, Sivagangai (Dt), Tamil Nadu, India, during the month of April 2017 and taxanomically identified by Dr. S. John Britto, Director, Rapinat Herbarium and Centre for Molecular Systematics (Authentication No. AR003 dated: 05/04/2017) at the St. Joseph's College (Campus), Tiruchirappalli, Tamil Nadu, India.

### **2.2 Extraction and fractionation**

The 3 kg freshly collected flowers were extracted with 90% ethanol. The combined alcoholic extract was concentrated *in vacuo* and the aqueous extract was successively fractionated with petroleum ether (60-80<sup>0</sup>C) (6x250 ml), Peroxide free diethyl ether (4x250 ml) and ethyl acetate (8x250 ml). Petroleum ether fraction and diethyl ether fraction did not give in any isolable compounds. Ethyl acetate fraction on concentration yielded a dry powder. The dried samples were dissolved in DMSO and were used for further studies.

## **2.3. IN VITRO ANTIOXIDANT ACTIVITY**

### **2.3.1 DPPH Assay method**

The DPPH free radical is reduced to a corresponding hydrazine, when it reacts with hydrogen donors. The DPPH radical is purple in colour and upon reaction with hydrogen donor changes to yellow colour. It is a decolouration assay, which is evaluated by the addition of the antioxidant to a DPPH solution in ethanol or methanol and the decrease in absorbance was measured at 490 nm <sup>[16]</sup>.

#### **Reagents:**

##### **A. Preparation of 2,2-Diphenyl 1-picryl hydrazyl solution (DPPH, 100 $\mu$ M):**

Twenty two milligrams of DPPH was accurately weighed and dissolved in 100 ml of methanol. From this stock solution, 18 ml was taken and diluted to 100ml using methanol to obtain 100  $\mu$ M DPPH solution.

##### **B. Preparation of test solutions:**

Twenty one milligrams of dry sample extracted from ethyl acetate was dissolved in distilled DMSO to get a solution of 21 mg/ml concentration. This solution was serially diluted to prepare lower concentrations.

##### **C. Preparation of standard solutions:**

Ten milligrams of ascorbic acid was accurately weighed and dissolved in 1 ml of Dimethyl sulfoxide (DMSO) to obtain 10 mg/ml concentrations. These solutions were serially diluted with DMSO to prepare lower concentrations.

#### **D. Procedure:**

The antioxidant activity was carried out in a 96 well micro titre plate. The final concentration of the dry sample extracted from ethyl acetate and standard solutions used 1000 µg/mL, 500 µg/mL, 125 µg/mL and 31.25 µg/mL were prepared. To 200µL of DPPH solution(violet colour), 10µL of each of the ethyl acetate fraction compound or the standard solution was added separately in wells of the micro titre plate. The plates were incubated at room temperature for 30 minutes. The absorbance of each solution was measured at 490 nm, using a micro plate reader. The antioxidants present in the test sample scavenge the DPPH free radicals and turn violet into yellow colour.

#### **2.3.2 ABTS free radical scavenging activity:**

ABTS free radical scavenging activity was carried out by 7.0 mM ABTS [2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid)] and 14.7 mM ammonium perdisulphate was dissolved in 5.0 ml demineralised water and stand 24 hrs at room temperature in the absence of sunlight, the blue green ABTS radical is formed in the solution. The solution was diluted then its absorbance is  $0.70 \pm 0.020$  at 734 nm is noted spectrophotometer<sup>[17]</sup>.

The different concentrations( 1000 µg/ml, 500 µg/ml, 125 µg/ml, 31.25 µg/ml) of the dry sample extracted from ethyl acetate fractions of *Moringa oleifera* flowers is prepared using ethanol as a solvent. 20.0 µL of the each sample solutions were added to 980.0 µL of ABTS radical solution and it was incubated in darkness for 10 minutes. The decrease in absorbance was noted at 734 nm. The anti-oxidant activity is determined by decolourization of the ABTS radical. 20.0 µL of ethanol containing a test tube was served as the control tube. Various concentrations of ascorbic acid (antioxidant) were used as reference compound.

### **2.4 ANTI- INFLAMMATORY ACTIVITY**

#### **2.4.1 HRBC membrane stabilization methodology:**

The anti-inflammatory activity of the dry sample extracted from ethyl acetate fractions of *Moringa oleifera* flowers was studied by HRBC stabilization method. The non-steroidal anti-inflammatory drugs are used to treat the inflammation<sup>[18]</sup>. The blood sample was collected from blood donors and confirmed who had not intake any Non-steroidal anti-inflammatory drugs for 2

weeks earlier to the experiment. The collected blood was mixed with equal quantity of Alsever solution (2 % dextrose, 0.8 % sodium citrate, 0.05 % citric acid and 0.42 % sodium chloride dissolved in water) and centrifuged at 3,000 rpm, then the packed cells washed with isosaline. 10 % suspension was prepared for the method. Various concentrations of dry sample extracted from ethyl acetate fractions of *Moringa oleifera* flowers were prepared in mg/ml using distilled water then added 1 ml of phosphate buffer, 2 ml hypo saline and 0.5 ml of HRBC suspension in each concentration and incubated at 37<sup>0</sup>C for 30 minutes. The mixture was centrifuged at 3,000 rpm for 20 minutes. The supernatant solution containing hemoglobin was predicted spectrophotometrically at 560 nm. Diclofenac (100 Jg/ml) was performed as a standard. The experiments were carried out in three sets and their mean values were used to calculate percentage (%) of HRBC membrane stabilization .<sup>[19]</sup>

**Percentage of Protection = (100- Optical Density of drug treated sample/ Optical Density of Control) X 100**

#### **2.4.2 Albumin denaturation methodology:**

The reaction mixture was containing dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers and 1% solution of bovine albumin fraction. A little amount of hydrochloric acid was added to the reaction mixture for adjusting the pH range.<sup>[20]</sup> The mixture was incubated at 37<sup>0</sup>C for twenty minutes. After the incubation process, the mixture was heated to 51<sup>0</sup>C for twenty minutes and cooled. The turbidity of the samples was measured spectrophotometrically at 660 nm. Diclofenac sodium was performed as a standard drug. The experiments were carried out in three sets and their mean values were used to calculate the inhibition percentage of protein denaturation was calculated.<sup>[21,22]</sup>

**Percentage of inhibition (%) =  
(Optical Density of Control- Optical Density of Sample/ Optical Density of Control) X 100**

### **3. RESULTS AND DISCUSSION**

#### **3.1 Anti-oxidant activity: DPPH assay**

The antioxidant activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers exhibited using 2,2-Diphenyl 1-picryl hydrazyl radical assay. In the present study, the free radical scavenging potentials of the extracts at different concentrations

were tested. This method determined the scavenging activity of DPPH radical. The activity of the sample is evidenced from the data presented in Table 1. The total antioxidant activity was increased with increasing concentrations of the dry sample extracted from ethyl acetate fraction is represented in figure 1. This result showed the IC<sub>50</sub> value (369.4 µg/ml) of the compound.

### **3.2 Anti-oxidant activity: ABTS assay**

ABTS radical assay also great technique for the investigation of antioxidant activity of antioxidants. The dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flower possesses effective antioxidant activity. It is evident from the data presented in Table 2. The result showed the percentage of cytotoxicity for various concentration of dry sample isolated from the ethyl acetate fraction of flowers of *Moringa oleifera*. The total antioxidant activity was gradually increased with abnormal increasing the concentrations of the dry sample extracted from ethyl acetate fraction is represented in figure 2. The IC<sub>50</sub> value of the assay was 211.9 µg/ml.

### **3.3 Anti-inflammatory activity of human red blood cell (HRBC) membrane stabilization method:**

The dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers exhibited significant anti-inflammatory activity of the human red blood cell (HRBC) membrane stabilization and the results are presented in Table 3. The results showed the percentage of inhibition in membrane stabilization gradually increased with the concentration of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers. The higher concentration 800 µg/ml showed significant stabilization ( $62.36 \pm 1.86$ ) towards human red blood cell membrane was represented in figure 3.

### **3.4 Anti-inflammatory activity of albumin denaturation method:**

Inflammation is caused by denaturation of proteins. The anti-inflammatory activity was investigated by albumin denaturation method. The inhibition of albumin denaturation activity exhibited by the various concentration of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers are given in Table 4. The results showed the effective inhibition in albumin denaturation. the inhibition activity was gradually increased with increasing the concentration of the test sample is represented in figure 4.

**Table 1: Anti-oxidant activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers by DPPH assay**

S. No	Concentration (µg/ml)	% Cytotoxicity (µg/ml)	IC <sub>50</sub> (µg/ml)
1	1000	70.54	369.4
2	500	63.01	
3	125	46.57	
4	31.25	29.45	

**Table 2 : Anti-oxidant activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers by ABTS assay.**

S. No	Concentration (µg/ml)	% Cytotoxicity(µg/ml)	IC <sub>50</sub> (µg/ml)
1	1000	76.34	211.9
2	500	68.19	
3	125	49.56	
4	31.25	36.78	


**Table 3: The human red blood cell membrane Stabilization activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers**

S. No	Concentration (µg/ml)	% of Inhibition
		Membrane Stabilization Mean ± S.E.M(S-I)
1	100	29.96 ± 0.41
2	200	39.48 ± 0.59
3	400	48.09 ± 0.61
4	600	58.93 ± 1.40
5	800	62.36 ± 1.86




**Table 4: The Inhibition of Albumin Denaturation activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers**


S. No	Concentration (µg/ml)	% of Inhibition
		Membrane Stabilization Mean ± S.E.M(S-I)
1	100	29.62 ± 0.58
2	200	39.18 ± 0.86
3	400	59.64 ± 0.94
4	600	68.17 ± 1.27
5	800	72.96 ± 1.49




**Figure1: Anti-oxidant activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers by DPPH assay**



**Figure 2: ABTS radical scavenging activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers**



**Figure 3: The human red blood cell membrane Stabilization activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers**



**Figure 4: The Inhibition of Albumin Denaturation activity of the dry sample extracted from ethyl acetate fraction of *Moringa oleifera* flowers**

#### **4.CONCLUSION**

In the present study, both DPPH assay and ABTS have showed a highest potential antioxidant activity and also the human red blood cell (HRBC) membrane stabilization activity

of the test sample. It could be concluded that the dry sample extracted from ethyl acetate fraction of flowers of *Moringa oleifera* is of phytopharmaceutical importance. The chemical compound present in the dry sample extracted from ethyl acetate fraction is ongoing in our research laboratory.

#### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

#### **ACKNOWLEDGEMENT**

The authors would like to thank Periyar E.V.R College, Trichy-23 and Alpha Omega biotech research centre, Salam.

#### **5. REFERENCES.**

1. Boonchum W, Peerapornpisal Y, Vacharapiyasophon P, Pekkoh J, Pumas C, Jamjai U, et al. Antioxidant activity of some seaweed from the gulf of Thailand. *Int J Agric Biol.* 2011; 13: 95–99.
2. Pakade V, Cukrowska E, Chimuka L. Comparison of antioxidant activity of *Moringa oleifera* and selected vegetables in South Africa. *South African Journal of Science.* 2013; 109: 1-5
3. Moon J, and Shibamoto T. Antioxidant assays for plant and food components. *J Agric Food Chem.* 2009; 57: 1655–1666.

4. Laandrault N, Pouchert P, Ravel P, Gase F, Cros G, Teissedro PL. Antioxidant activities and phenolic level of French wines from different varieties and vintages. *J Agric Food Chem.* 2001; 49: 3341–3343.
5. Mukazayire MJ, Allaeyes V, Duez P, Calderon PB, Stevigny C, and Bigendako MJ. Evaluation of the hepatotoxic and hepatoprotective effect of Rwandese herbal drugs on in vivo (guinea pigs barbiturate-induced sleeping time) and in vitro (rat precision-cut liver slices, PCLS) models. *Experimental and Toxicologic Pathology.* 2010; 62(3): 289–299.
6. Marletta, M.A. Nitric oxide synthases structure and mechanism. *J. Biol. Chem.* 1993; 268: 12231–12234.
7. Watkins LR, Milligan ED, Maier SF, Glial proinflammatory cytokines mediate exaggerated pain states: implications for clinical pain. *Advances in Experimental Medicine and Biology.* 2003; 521: 1–21.
8. Smejkal K, Selected natural phenolic compounds - potential treatment for peripheral neuropathy?. *Ceska Slovenska Farmacie.* 2014; 63:55–70.
9. Nadkarni K.M, Nadkarni A. K and Chopra R.N. *The Indian Materia Medica.* Popular Prakasan, Bombay. 1954; 1: 811 – 816.
10. Fahey JW. *Moringa oleifera*: A review of the medical evidence for its nutritional, therapeutic, and prophylactic properties. Part 1. *Tree for Life Journal.* 2005; 1:5-20.
11. Torres-Castillo JA, Sinagawa-Garcia SR, Martinez-Avila GCG, Lopez-Flores AB, Sanchez-Gonzalez EI, Aguirre-Arzola VE, Torres-Acosta RI, Olivares-Saenz E, Osorio-Hernandez E and Gutierrez-Diez A. *Moringa oleifera* phytochemical detection, antioxidants, enzymes and antifungal Properties. *Phyton International journal of experimental botony.* 2013; 82: 193-202.

12. Anwar F, Latif S, Ashraf M and Gilani A.H. *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytotherapy Research*. 2007; 21: 17-25.
13. Goyal, B.R., Agrawal B.B, Goyal R.K and Mehta A.A. Phyto- pharmacology of *Moringa oleifera* Lam. An overview. *Natural Product Radiance*. 2007; 4: 347-353.
14. Chumark P, Khunawat Y, Sanvarinda S, Phornchirasilp N, Phumala Morales L, Phivthongngam P, Ratanachamnong S, Srisawat and Pongrapeeporn K.S. The in vitro and ex vivo antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera* Lam. leaves. *Journal of Ethnopharmacology*. 2008; 116: 439-446.
15. Peixoto J.R, Silva G.C, Costa R.A, De Sousa Fontenell J.R, Vieira G.H, Filho A.A, and Dos Fernandes Vieira R.H. In vitro antibacterial effect of aqueous and ethanolic *Moringa* leaf extracts. *Asian Pacific Journal of Tropical Medicine*. 2011; 4: 201-204.
16. Ilavarsan R, Mallika M, Venkataraman S. Anti-inflammatory and antioxidant activities of cassia fistula linn bark extracts. *African Journal of Traditional, Complementary and Alternative Medicines*. 2005; 2(1):70-85.
17. Roberta Re, Nicoletta Pellegrini, Anna Proteggente, Ananth Pannala, Min Yang and Catherine Rice-Evans. Antioxidant Activity Applying An Improved Abts Radical Cation Decolorization Assay. *Free Radical Biology & Medicine*. 1999; 26: 1231–1237.
18. Gopalakrishnan S, Vadivel E and Dhanalakshmi K. Anti-inflammatory and analgesic activities of *Tephrosia purpurea* Linn. aerial and root extracts. *Journal of Pharmacy Research* 2010; 3: 1103-1106.
19. Nagaharika Y, kalyani V, Rasheed S and karthikeyan R. Anti inflammatory activity of leaves of *Jatropha gossypifolia* L. by hrbc membrane stabilization method. *Journal of Acute Disease*. 2013; 156-158

20. Sakat S, Juvekar AR, Gambhire MN. In vitro antioxidant and inflammatory activity of methanol extract of *Oxalis corniculata* Linn. *International Journal of Pharma and Pharmacological Sciences*. 2010; 146-155.

21. Brown JH and Mackey HK. Inhibition of heat-induced denaturation of serum proteins by mixtures of nonsteroidal anti-inflammatory agents and aminoacids. *Proceedings of the Society for Experimental Biology and Medicine*. 1968; 128: 225-8.

22. Solomon S, Murugananthan N and Senthamilselvi M.M. Anti-oxidant and anti-inflammatory activity of *Thespesia populnea* (flowers). *Pharmacophore*. 2015; 6(1): 53-59.