

**ECONOMICAL EFFICACY OF WEED MANAGEMENT OPTIONS IN  
GROUNDNUT + PIGEONPEA RELAY INTERCROPPING SYSTEM**

**Abstract**

- **Aims:** To evaluate the effect of different weed management options on the economics in groundnut + pigeonpea relay intercropping on medium black clayey soils of Junagadh .
- **Study design:** Field experiment was conducted at Junagadh during *kharif* 2019-20 and 2020-21 in Randomized Block Design with three replications to evaluate the effect of different weed management options on the economics in groundnut + pigeonpea relay intercropping on medium black clayey soils. The treatments comprised of: pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* interculturing and hand weeding at 45 DAS, pendimethalin 0.45 kg ha<sup>-1</sup> + oxyfluorfen 0.09 kg ha<sup>-1</sup> as PE *fb* interculturing and hand weeding at 45 DAS), interculturing and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as POE, interculturing and hand weeding at 15 DAS *fb* quizalofop p ethyl 40 g ha<sup>-1</sup> at 45 DAS as POE, interculturing and hand weeding at 15 DAS *fb* propaquizafop 70 g ha<sup>-1</sup> at 45 DAS as POE, pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as POE, pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* quizalofop p ethyl 40 g ha<sup>-1</sup> at 45 DAS as POE, pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* propaquizafop 70 g ha<sup>-1</sup> at 45 DAS as POE (T<sub>8</sub>), weed free, and unweeded control (T<sub>10</sub>).
- **Place and duration of the study :** Instructional Farm, Department of Agronomy, College of Agriculture, Junagadh Agricultural University, Junagadh during *kharif* seasons of 2019-20 and 2020-21.
- **Methodology:** on the basis of prevailing local charges. The gross realization in terms of rupees per hectare. A net return of each treatment was calculated by deducting the total cost of cultivation from the gross returns.
- **Result:** Significantly higher amount of gross returns were recorded with the weed free treatment (T<sub>9</sub>), which was closely followed by interculturing and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE (T<sub>3</sub>). Statistically higher net returns and higher net

returns and B:C ratio were registered with interculture and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE (T<sub>3</sub>).

*Key words: Groundnut, pigeonpea, interculturing, pendimethalin, oxyfluorfen, sodium acifluorfen, clodinafop propargyl, quizalofop-p-ethyl and propaquizafop*

## Introduction

Inadequate and erratic rainfall along with weed, insect and other pest attacks prove to be the reason for lower yields and in extreme cases may result in complete failure of the crop. There is a greater difference between the demand and supply of edible oil due to low productivity of oilseed crops. Furthermore, the shortage of pulses and oilseeds has also intensified the problem of malnutrition. Thus, practice of intercropping system offers scope for maximizing and stabilizing returns from *kharif* oilseed crops rather than as a sole crop. In order to reap higher benefits from cultivation of pulses and oilseed crops, there is an urgent need to increase the area and the productivity of these crops. Due to the limiting factors like industrialization and urbanization, the scope for expansion of area under pulses and oilseeds is minimal. This situation instigates the need to adopt appropriate agronomic measures required to increase production (Chaudari *et al.*, 2017). Intercropping has been in vogue for long time to sustain yield, minimize risk, utilize the lag phase and improve productivity. Selection of an appropriate and compatible component crops for intercropping system is necessary for reducing plant competition for resources (Singh *et al.*, 2009). To stabilize crop production and to provide insurance against aberrant weather situation, relay intercropping could be a viable agronomic option for risk minimizing, more profit and sustainable venture. It provides an opportunity to use per unit land, water, nutrient and money invested efficiently. Substantial yield advantage can be achieved through inter/relay cropping as compared to sole cropping. Groundnut + pigeonpea is an emerging relay intercropping system in India, especially in Gujarat and Maharashtra ([www.jau.in](http://www.jau.in)).

The current shortage of both pulse and vegetable oils in India has stimulated thought on developing new systems of pigeonpea and groundnut intercropping. The approach may involve growing of a long duration crop like pigeonpea with groundnut, so that the loss sustained if any, groundnut may be compensated with long duration crop (Gunri *et al.*, 2015). Intercropping of groundnut + pigeonpea system in 2:1 row

proportion is the recommended row proportion for southern Saurashtra region of Gujarat especially when semi-spreading groundnut varieties (GG-20) are used as recommended by Main Oilseeds Research Station, JAU, Junagadh. This combination is particularly prevalent on red soils of the Southern states of India.

Weed infestation is one of the major constraints in productivity of any crop. The slow initial growth of groundnut favours the weed growth and reduces yield up to 75% (Gnanamurthy and Balasubramaniyan, 1998). Adoption of manual weeding though efficient but costly too. Further availability of labour at appropriate time is another constraint which enables the vigorous growth of weeds to compete at initial stages. Use of herbicides could be an alternative and economically feasible method of weed control under these conditions. However, the success of weed control could be determined by the choice of suitable and safe herbicides for both sole and intercrop (Nambi *et al.*, 2006). In the present study, the pigeonpea crop is planned to be sown one month after the sowing of groundnut due to the vigorous growth of pigeonpea may affect the groundnut during the crucial periods like flowering *etc.* Also, the different dates of sowing reduce the competition between the component crops and thereby increasing the yield of the crops. So, for the further increase in productivity of both the crops efficient weed control practices are needed as both the crops have slower initial growth. For the efficient weed management in the cropping system, pre-emergence as well as post-emergence herbicides are included to provide greater advantage to the crops reducing the competition effect of the weeds. The crop production per unit land can be increased with appropriate cropping system and weed management which also must be an economical feasible method which can be adopted by the farmers.

### **Materials and Methods**

The study was conducted at Instructional Farm, Department of Agronomy, College of Agriculture, Junagadh Agricultural University, Junagadh during *kharif* seasons of 2019-20 and 2020-21. Geographically, Junagadh is located at 21.5° N latitude and 70.5° E longitude with an altitude of 60 m above the mean sea level on the western side on the foothill of mountain 'Girnar' under South Saurashtra Agro-climatic Zone of Gujarat state and enjoys a typically subtropical climate characterized by fairly cold and dry winter, hot and dry summer and warm and moderately humid monsoon.

The soil of the experimental plot was clayey in texture, medium in organic carbon (0.62 %), slightly alkaline in reaction with pH (8.32) and EC (0.286 dS m<sup>-1</sup>) in 2019-20. The soil was medium in available nitrogen (270.20 kg ha<sup>-1</sup>), medium in available phosphorus (28.2 kg ha<sup>-1</sup>) and medium in available potash (251.7 kg ha<sup>-1</sup>) in 2019-20. In the study, additive series of intercropping was adopted, wherein the main/base crop is groundnut and intercrop is pigeonpea. “Gujarat Groundnut 20” variety and pigeonpea variety “Gujarat Junagadh Pigeonpea 1” as intercrop. to carry out the present investigation. The treatments comprised of pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* interculturing and hand weeding at 45 (DAS) (T<sub>1</sub>), pendimethalin 0.45 kg ha<sup>-1</sup> + oxyfluorfen 0.09 kg ha<sup>-1</sup> as PE *fb* interculturing and hand weeding at 45 DAS (T<sub>2</sub>), interculturing and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as POE (T<sub>3</sub>), interculturing and hand weeding at 15 DAS *fb* quizalofop p ethyl 40 g ha<sup>-1</sup> at 45 DAS as POE (T<sub>4</sub>), interculturing and hand weeding at 15 DAS *fb* propaquizafop 70 g ha<sup>-1</sup> at 45 DAS as POE (T<sub>5</sub>), pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as POE (T<sub>6</sub>), pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* quizalofop p ethyl 40 g ha<sup>-1</sup> at 45 DAS as POE (T<sub>7</sub>), pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* propaquizafop 70 g ha<sup>-1</sup> at 45 DAS as POE (T<sub>8</sub>), weed free (T<sub>9</sub>), and unweeded control (T<sub>10</sub>).

### 2.3 Experimental design

The objective of experiment was to to evaluate the effect of different weed management options on the economics in groundnut + pigeonpea relay intercropping on medium black clayey soils of Junagadh. The experiment was laid out in randomized block design with ten treatments, which are replicated thrice.

As the research proposal is on weed management, the practices were adopted in accordance to the proposed treatments. Interculturing in combination with hand weeding was carried in treatments 3, 4 and 5 at 15 DAS while in treatment 1 and 2 it was conducted at 45 days after sowing (DAS) of groundnut. Pre emergence herbicide *viz.*, pendimethalin 30% EC and oxyfluorfen 23.5% EC were applied on the next day of sowing of groundnut and post emergence herbicides, quizalofop-p-ethyl, propaquizafop and sodium acifluorfen + clodinafop propargyl were applied at 45 DAS after intercultivation. The weed free was maintained clean with regular manual

weeding. The unweeded control was left **unweeding** allowing continuous growth of weeds.

**Economics of Treatments:** The expenses incurred for all the cultivation operations from preparatory tillage to harvesting including the cost of inputs *viz.*, seeds, manures, irrigation, biopesticides, *etc.* applied to each treatment was calculated on the basis of prevailing local charges. The gross realization in terms of rupees per hectare was worked out taking into consideration the pod and haulm yields of groundnut and seed and stalk yields of pigeonpea from each treatment and local market prices. A net return of each treatment was calculated by deducting the total cost of cultivation from the gross returns. The Benefit : Cost (B:C) ratio was calculated with the help of following formula.

$$\text{B:C ratio} = \frac{\text{Gross returns } (\text{₹ ha}^{-1})}{\text{Total cost of cultivation } (\text{₹ ha}^{-1})}$$

## Results

The data on economics of weed management treatments in groundnut + pigeonpea relay intercropping system during individual years (2019-20 and 2020-21) as well as average of both the **years furnished** in Table 1-4. Net returns and cost of cultivation graphically portrayed in Fig. 1.

### Cost of cultivation

**Among** the different weed management treatments, the weed free treatment (T<sub>9</sub>) had higher cost of cultivation (₹ 74424, 75584 and 75004 ha<sup>-1</sup> during 2019-20, 2020-21 as well as pooled results, respectively), followed by the treatments T<sub>3</sub> (**interculturing** and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (**ready** mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE), T<sub>6</sub> (**pendimethalin** 0.9 kg ha<sup>-1</sup> as aPE *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE), T<sub>2</sub> (**pendimethalin** 0.45 kg ha<sup>-1</sup> + oxyfluorfen 0.09 kg ha<sup>-1</sup> as PE *fb* interculturing and hand weeding at 45 DAS), T<sub>1</sub> (**pendimethalin** 0.9 kg ha<sup>-1</sup> as PE *fb* interculturing and hand weeding at 45 DAS), T<sub>5</sub> (**interculturing** and hand weeding at 15 DAS *fb* propaquizafop 70 g ha<sup>-1</sup> at 45 DAS as PoE), T<sub>4</sub> (**interculturing** and hand weeding at 15 DAS *fb* quizalofop-p-ethyl 40 g ha<sup>-1</sup> at 45 DAS as PoE), T<sub>8</sub> (**pendimethalin** 0.9 kg ha<sup>-1</sup> as PE *fb* propaquizafop 70 g ha<sup>-1</sup> at 45 DAS as PoE), T<sub>7</sub> (**pendimethalin** 0.9 kg ha<sup>-1</sup> as PE *fb* quizalofop-p-ethyl 40 g ha<sup>-1</sup> at 45 DAS as

PoE). The lowest cost of cultivation ( $\square$  49277, 50437 and 49857 ha<sup>-1</sup> during 2019-20, 2020-21 as well as pooled results respectively) was recorded with the unweeded control (T<sub>10</sub>) during 2019-20, 2020-21 as well as pooled results

### Gross Returns

Significantly gross returns ( $\square$  231265, 208047 and 219657 /ha during 2019-20, 2020-21 as well as pooled results respectively) were recorded with the weed free treatment (T<sub>9</sub>) during 2019-20, 2020-21 and pooled results. During 2019-20, 2020-21, the weed free treatment (T<sub>9</sub>) statistically at par with interculturing and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE (T<sub>3</sub>) and pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE (T<sub>6</sub>). While in pooled results it was observed that T<sub>9</sub> was on par with T<sub>3</sub> alone. Lower gross returns ( $\square$  65215, 50530 and 57872 /ha during 2019-20, 2020-21 and pooled results, respectively) were obtained with the unweeded control.

### 4.5.3 Net Returns

Statistically higher net returns ( $\square$  161201, 133584 and 147393 /ha during 2019-20, 2020-21 as well as pooled results respectively) were registered with interculturing and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE (T<sub>3</sub>) during 2019-20, 2020-21 and pooled results at par with weed free treatment (T<sub>9</sub>) and pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE (T<sub>6</sub>) during 2019-20, 2020-21 and pooled data. The lower net returns ( $\square$  15938, 95 and 8015 /ha during 2019-20, 2020-21 and pooled results, respectively) were obtained with the unweeded control (T<sub>10</sub>).

### 4.5.4 B:C ratio

Among weed management treatments, the highest B:C ratio (3.84, 3.30 and 3.57 during 2019-20, 2020-21 as well as pooled results, respectively) was obtained with T<sub>3</sub> (interculturing and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE), followed by the treatments T<sub>6</sub> (pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (ready mix) 1 kg ha<sup>-1</sup> at 45 DAS as PoE), T<sub>1</sub> (pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* interculturing and hand weeding at 45 DAS), T<sub>2</sub> (pendimethalin

0.45 kg ha<sup>-1</sup> + oxyfluorfen 0.09 kg ha<sup>-1</sup> as PE *fb* interculturing and hand weeding at 45 DAS), T<sub>9</sub> (weed free treatment), T<sub>4</sub> (interculturing and hand weeding at 15 DAS *fb* quizalofop-p-ethyl 40 g ha<sup>-1</sup> at 45 DAS as PoE), T<sub>5</sub> (interculturing and hand weeding at 15 DAS *fb* propaquizafop 70 g ha<sup>-1</sup> at 45 DAS as PoE), T<sub>7</sub> (pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* quizalofop-p-ethyl 40 g ha<sup>-1</sup> at 45 DAS as PoE) and T<sub>8</sub> (pendimethalin 0.9 kg ha<sup>-1</sup> as PE *fb* propaquizafop 70 g ha<sup>-1</sup> at 45 DAS as PoE). The treatment of unweeded control (T<sub>10</sub>) recorded the lowest B:C ratio (1.32, 1.00 and 1.16 during 2019-20, 2020-21 and pooled results, respectively) during 2019-20, 2020-21 along with pooled results.

## Discussion

Higher net returns and B:C ratio were registered eventhough, the expenditure was higher on weed management in treatments T<sub>3</sub>, it was compensated by increased economic yield of groundnut and pigeonpea with higher gross returns. Consequently, higher B:C ratio was obtained. The unweeded control (T<sub>10</sub>) registered lower yields of groundnut and pigeonpea, thereby resulted in lower gross returns and net returns, even though the cost of cultivation was lower and finally leading to lower B:C ratio. Similar results were also reported by Shinde *et al.* (2003) in pigeonpea + pearl millet intercropping system under integrated weed management system. The results are in line with those of Bundhar and Tamilselvan (2003), Sasikala *et al.* (2004), Solanki *et al.* (2005), Chandolia *et al.* (2010), Malunjar *et al.* (2012), Padmaja *et al.* (2013), Rai *et al.* (2016), Reddy *et al.* (2016) and Priya *et al.* (2017).

## Conclusion

The evaluation of the economics indicated that higher net return and B:C ratio were recorded with interculturing and hand weeding at 15 DAS *fb* sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha<sup>-1</sup> at 45 DAS

## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## References

- Bundhar M N and Tamilselvan N T 2003. Weed dynamics, yield and economics in sorghum (*Sorghum bicolor*) + pulses intercropping system under rainfed condition. *Indian Journal of Agronomy* **48**(2) : 93-95.
- Chandolia P C, Dadheech R C Solanki N S and Mundra S L 2010. Weed management in groundnut (*Arachis hypogaea* L.) under varying crop geometry. *Indian Journal of Weed Science* **42**(3&4) : 235-237.
- Chaudari D T, Vekariya P D, Vora V D, Talpada M M and Sutaria S G 2017. Enhancing productivity of groundnut based intercropping systems under rainfed conditions of Gujarat. *Legume Research* **40**(3) : 520-525.
- Gnanamurthy, P. and Balasubramaniyan, P. 1998. Weed management practices and their influence on weed growth and yield of groundnut. *Indian Journal of Agronomy*, **43**: 122-125.
- Gunri S K, Nath R and Puste A M 2015. Effect of plant density and fertility levels of *kharif* groundnut (*Arachis hypogaea* L.) + pigeon pea (*Cajanus cajan*) intercropping system under red and laterite zone of West Bengal, India. *Legume Research* **38**(5) : 639-642.
- <http://www.jau.in/index.php/2-init-content/research/101-pulses-research-station>
- Malunekar B D, Mulik B B and Patil S C 2012. Evaluation of post-emergence herbicides in rainy season groundnut. *Indian Journal of Weed Science* **44**(2) : 95-97.
- Nambi J, Sundari A and Pandian B J 2006. Weed management in groundnut-based intercropping system. *Indian Journal of Weed Science* **38**(1 & 2) : 159-160.
- Padmaja B, Reddy M M and Reddy D V V 2013. Weed control efficiency of pre- and post-emergence herbicides in pigeonpea (*Cajanus cajan* L.). *Journal of Food Legumes* **26** : 44-45.
- Priya R, Chinnusamy C, Murali P A, Janaki P and Babu C 2017. Evaluation of oxyfluorfen (23.5% EC) herbicide on weed control, economics and profitability of groundnut in the western zone of Tamil Nadu. *Chemical Science Review and Letters* **6**(21) : 88-93.
- Rai C L, Tiwari R K, Sirothia P and Pandey S 2016. Intercropping and weed management effects on weed dynamics, productivity and economics of pigeonpea. *Indian Journal of Weed Science* **48**(1) : 44-47.
- Reddy N, Vidyasagar G E Ch. and Laxminarayana P 2016. Integrated weed management in *rabi* groundnut (*Arachis hypogaea* L.). *International Journal of Current Research* **8**(11) : 40883-40885.
- Sasikala B, Reddy Y and Reddy R C 2004. Pre- and postemergence herbicides on weed control and yield of groundnut (*Arachis hypogaea*). *Indian Journal of Dryland Agriculture Research and Development* **19**(1) : 78-80.
- Shinde H S, Pawar V S, Suryawanshi G B, Ahire N R and Surve U S 2003. Integrated weed management studies in pigeonpea + pearl millet intercropping (2: 2) system. *Indian Journal of Weed Science*, **35** : 90-92.
- Singh K K, Ali M and Venkatesh M S 2009. Pulses in cropping system. *Technical Bulletin*, IIPR, Kanpur.
- Solanki R M, Bhalu V B, Jadav K V and Kelaiya G R 2005. Studies on integrated weed management in irrigated groundnut. *Indian Journal of Weed Science* **37**(1&2) : 119-120.
- Willey R W 1979. Intercropping its importance and research needs, *Agronomy of research approaches*. *Field Crop Abstract* **32** : 1-10.



**Table-1: Cost of cultivation of various weed management practices in groundnut + pigeonpea relay intercropping system**

Treatment	Cost of cultivation (₹ ha <sup>-1</sup> )		
	2019	2020	Pooled
T <sub>1</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> interculturing and hand weeding at 45 DAS.	53909	55069	54489
T <sub>2</sub> :Pendimethalin 0.45 kg ha <sup>-1</sup> + oxyfluorfen 0.09 kg ha <sup>-1</sup> PE <i>fb</i> interculturing and hand weeding at 45 DAS.	53978	55138	54558
T <sub>3</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha <sup>-1</sup> at 45 DAS as PoE.	56853	58014	57434
T <sub>4</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> quizalofop-p-ethyl 40 g ha <sup>-1</sup> at 45 DAS as PoE.	53675	54835	54255
T <sub>5</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> propaquizafop 70 g ha <sup>-1</sup> at 45 DAS as PoE.	53786	54947	54367
T <sub>6</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha <sup>-1</sup> at 45 DAS as PoE.	55897	57057	56477
T <sub>7</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> quizalofop-p-ethyl 40 g ha <sup>-1</sup> at 45 DAS as PoE.	52718	53878	53298
T <sub>8</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> propaquizafop 70 g ha <sup>-1</sup> at 45 DAS as PoE.	52830	53990	53410
T <sub>9</sub> :Weed free	74424	75584	75004
T <sub>10</sub> :Unweeded control	49277	50437	49857

**Table-2: Gross returns of various weed management practices in groundnut + pigeonpea relay intercropping system**

Treatment	Gross returns (□ ha <sup>-1</sup> )		
	2019	2020	Pooled
T <sub>1</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> interculturing and hand weeding at 45 DAS.	190075	171022	180548
T <sub>2</sub> :Pendimethalin 0.45 kg ha <sup>-1</sup> + oxyfluorfen 0.09 kg ha <sup>-1</sup> PE <i>fb</i> interculturing and hand weeding at 45 DAS.	187761	158451	173106
T <sub>3</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha <sup>-1</sup> at 45 DAS as PoE.	218055	191598	204826
T <sub>4</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> quizalofop-p-ethyl 40 g ha <sup>-1</sup> at 45 DAS as PoE.	164664	134586	149625
T <sub>5</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> propaquizafop 70 g ha <sup>-1</sup> at 45 DAS as PoE.	156506	130665	143585
T <sub>6</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha <sup>-1</sup> at 45 DAS as PoE.	206900	181295	194098
T <sub>7</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> quizalofop-p-ethyl 40 g ha <sup>-1</sup> at 45 DAS as PoE.	146853	120346	133599
T <sub>8</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> propaquizafop 70 g ha <sup>-1</sup> at 45 DAS as PoE.	141433	110037	125735
T <sub>9</sub> :Weed free	231265	208047	219656
T <sub>10</sub> :Unweeded control	65215	50530	57872
S.Em.±	7075	6354	4755
C.D. at 5%	21021	18878	13637
C.V. %	10.64	12.40	11.42

**Table-3: Net returns of various weed management practices in groundnut + pigeonpea relay intercropping system**

Treatment	Net returns (₹ ha <sup>-1</sup> )		
	2019	2020	Pooled
T <sub>1</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> interculturing and hand weeding at 45 DAS.	136166	115953	126060
T <sub>2</sub> :Pendimethalin 0.45 kg ha <sup>-1</sup> + oxyfluorfen 0.09 kg ha <sup>-1</sup> PE <i>fb</i> interculturing and hand weeding at 45 DAS.	133783	103313	118548
T <sub>3</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha <sup>-1</sup> at 45 DAS as PoE.	161201	133584	147393
T <sub>4</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> quizalofop-p-ethyl 40 g ha <sup>-1</sup> at 45 DAS as PoE.	110989	79751	95370
T <sub>5</sub> :Interculturing and hand weeding at 15 DAS <i>fb</i> propaquizafop 70 g ha <sup>-1</sup> at 45 DAS as PoE.	102720	75718	89219
T <sub>6</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha <sup>-1</sup> at 45 DAS as PoE.	151003	124238	137621
T <sub>7</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> quizalofop-p-ethyl 40 g ha <sup>-1</sup> at 45 DAS as PoE.	94134	66468	80301
T <sub>8</sub> :Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> propaquizafop 70 g ha <sup>-1</sup> at 45 DAS as PoE.	88604	56047	72325
T <sub>9</sub> :Weed free	156840	132463	144652
T <sub>10</sub> :Unweeded control	15938	92	8015
S.Em.±	7075	6354	4755
C.D. at 5%	21021	18878	13637
C.V. %	7.17	7.56	7.36

**Table-4: Benefit: cost ratio of various weed management practices in groundnut + pigeonpea relay intercropping system**

Treatment	B:C ratio		
	2019	2020	Pooled
T <sub>1</sub> : Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> interculturing and hand weeding at 45 DAS.	3.53	3.11	3.32
T <sub>2</sub> : Pendimethalin 0.45 kg ha <sup>-1</sup> + oxyfluorfen 0.09 kg ha <sup>-1</sup> PE <i>fb</i> interculturing and hand weeding at 45 DAS.	3.48	2.87	3.18
T <sub>3</sub> : Interculturing and hand weeding at 15 DAS <i>fb</i> sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha <sup>-1</sup> at 45 DAS as PoE.	3.84	3.30	3.57
T <sub>4</sub> : Interculturing and hand weeding at 15 DAS <i>fb</i> quizalofop-p-ethyl 40 g ha <sup>-1</sup> at 45 DAS as PoE.	3.07	2.45	2.76
T <sub>5</sub> : Interculturing and hand weeding at 15 DAS <i>fb</i> propaquizafop 70 g ha <sup>-1</sup> at 45 DAS as PoE.	2.91	2.38	2.64
T <sub>6</sub> : Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> sodium acifluorfen 16.5% + clodinafop propargyl 8% (Ready mix) 1 kg ha <sup>-1</sup> at 45 DAS as PoE.	3.70	3.18	3.44
T <sub>7</sub> : Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> quizalofop-p-ethyl 40 g ha <sup>-1</sup> at 45 DAS as PoE.	2.79	2.23	2.51
T <sub>8</sub> : Pendimethalin 0.9 kg ha <sup>-1</sup> PE <i>fb</i> propaquizafop 70 g ha <sup>-1</sup> at 45 DAS as PoE.	2.68	2.04	2.36
T <sub>9</sub> : Weed free	3.11	2.75	2.93
T <sub>10</sub> : Unweeded control	1.32	1.00	1.16




Fig-1: Economics of weed management treatments

UNDER PEER REVIEW