

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11

# Impact of zinc fertilization on yield, yield attributes and quality parameters of finger millet varieties under rainfed *alfisols* of southern zone, Andhra Pradesh

12  
13

---

## ABSTRACT

**Aims:** To study the effect of zinc fertilization through soil and foliar at different stages of finger millet on yield, yield attributes and quality in two major finger millet varieties under rainfed *alfisols* of southern zone, Andhra Pradesh

**Study design:** Split-plot design

**Place and Duration of Study:** Wetland farm, S.V Agricultural College, Tirupati and during *kharif* season of 2019 and 2020 (Two seasons)

**Methodology:** Zinc fertilization to two major finger millet varieties *viz.*, Vakula and Tirumala through soil and foliar application at different crop stages with following treatments *viz.*, Control (No fertilizers and manures); RDF (60 -30-20 kg N-P-K + FYM @ 4 t ha<sup>-1</sup>); RDF + soil application of ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> as basal; RDF+Soil application of chelated-ZnSO<sub>4</sub> @ 5 kg ha<sup>-1</sup>; RDF+foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence stage; RDF+foliar application of 0.2% ZnSO<sub>4</sub> at grain filling stage; and RDF+foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence and grain filling stage. The yield, yield attributes and quality parameters *viz.*, protein, zinc and iron content in grains were determined by adopting standard protocols.

**Results:** The application of zinc significantly ( $p \leq 0.05$ ) improved the yield and quality parameters over control. The foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence and grain filling stage was significantly ( $p \leq 0.05$ ) improved the yield and yield attributes of finger millet over RDF. The grain yield, straw yield, no. of productive tillers per plant, no. of fingers per plant were increased to 57.0%, 83.2%, 44.6% and 51.7%, respectively over RDF *i.e.*, 60-30-20 kg N-P-K + FYM @ 4 t ha<sup>-1</sup>. The quality parameters namely protein, grain zinc and iron also increased up to 40.7%, 69.5% and 43.2%, respectively over RDF.

**Conclusion:** Application of zinc sulphate at ear head emergence and grain filling stages enhanced the yield, yield parameters and quality parameters compared to other treatments of tirumala variety under rainfed *alfisols* of southern zone of Andhra Pradesh.

14  
15  
16  
17

*Keywords:* Zn fertilization; fortification; grain yield; protein, Recommended fertilizer dose

## 1. INTRODUCTION

18  
19  
20  
21  
22  
23  
24  
25

Finger millet (*Eleusine coracana*) is commonly known as “Nutritious millet” owing to its nutritional superiority over many cereal crops (rice, maize and sorghum) in terms of proteins, minerals, iron, calcium and vitamins. The grains contains about 5 to 8 per cent protein, 65 to 75 per cent carbohydrates, 15 to 20 per cent dietary fiber and 2.5 to 3.5 per cent minerals and it has 30 times more calcium than rice (344 mg/100 g). Finger millet well recognized with their health beneficial effects *viz.*, anti-diabetic, anti-tumorigenic, atherosclerogenic, antioxidant and antimicrobial properties (1).

26 Finger millet extensively cultivated in the tropical and sub-tropical regions, which  
27 accounts for about 85% of total millet production in India. It is an important small millet crop  
28 ranked third in cultivated area, production and productivity of 1.19 mha, 1.98 mt and 1661 kg  
29 ha<sup>-1</sup>, respectively (2) and it has the pride of place in having the highest productivity among  
30 the millets after sorghum and pearl millet (3). In Andhra Pradesh it covers an area of 31.63  
31 thousand ha with a production and productivity of 35,000 tonnes and 1087 kg ha<sup>-1</sup>,  
32 respectively. The grains have long storability even under ambient conditions and have made  
33 them “famine reserves”. This aspect is at most important as Indian agriculture suffers from  
34 vagaries of monsoon (4). Under increased probability of occurrence of drought and soil  
35 fertility degradation, many farmers opted to raise this crop, hence the cultivated area was  
36 allocated for this crop has significantly ( $p < 0.05$ ) increased over the last decade (5).

37 Zinc (Zn) is considered the major limiting micronutrient in most of the areas limiting  
38 the crop yields. Zinc has role in diverse physiological functions in biological systems. Zinc is  
39 typically the second most abundant transition metal in organisms after iron and the only  
40 metal represented in all six enzyme classes viz., oxidoreductase, transferases, hydrolases,  
41 lyases, isomerases and ligases (6). Further around 200 enzymes responsible for growth,  
42 development, immune function and resistance to infections are regulated by zinc in plant  
43 system (7). Hence, Zinc insufficiency in soils may cause lower yield or sometimes crop  
44 failure and leads to poor accumulation of zinc into grains causing zinc malnutrition in  
45 humans (8) (9). Zinc deficiency is 5<sup>th</sup> leading cause of deaths in the developing world and  
46 about 0.8 million people die annually of which 0.45 million are children under the age of five  
47 as per WHO reports. The extent of zinc deficiency was 49% in Indian soils (10). The soils of  
48 Andhra Pradesh are also deficient in Zn, hence there is a dietary need to increase grain Zn  
49 content of finger millet, improving the remobilization of absorbed and accumulated Zn to  
50 grain is a research priority. Zn application also reported to increase the grain iron (Fe)  
51 concentration in pearl millet (11).

52 The agronomic bio-fortification is a easier and faster approach to increase grain Zn  
53 concentration in finger millet. Several studies revealed that Zn fertilization increased Zn  
54 concentration in rice grain from 35 to 141 percent (12), increased from 24 to 48 percent in  
55 wheat (13) and up to 72 percent in maize (14). But limited research has done on impact of  
56 zinc fertilization on yield, quality and bio-fortification in finger millet. Hence present study was  
57 initiated to study the effect of Zn fertilization on yield, yield attributes and grain fortification in  
58 finger millet under rainfed condition.

## 60 2. MATERIAL AND METHODS

### 62 2.1 Description of experimental site

#### 64 2.1.1 Climate

66 . Field experiment was carried out at wetland farm, S.V.Agricultural College, Tirupati  
67 Andhra Pradesh, India during *kharif* season in the year 2019 and 2020. Geographically  
68 located between 13.5° N and 79.5° E with an altitude of 182.9 m above MSL. The region has  
69 a semi-arid type climate. During *kharif* 2019, crop received 769.0 mm of rainfall in 43 rainy  
70 days with standard week wise mean maximum and minimum temperature ranged from 28.2  
71 to 36.1°C and 21.3 to 27.6°C, respectively. The mean sunshine hours and mean evaporation  
72 (USWB Class-A Open Pan evaporimeter) ranged from 2.0 to 8.8 hours day<sup>-1</sup> and 1.9 to 6.0  
73 mm per day with an average of 4.7 hr day<sup>-1</sup> and 4.2 mm per day, respectively. During *kharif*  
74 2020, crop received 723.8.0 mm of rainfall in 31 rainy days with standard week wise mean  
75 maximum and minimum temperature ranged from 28.4 to 36.0°C and 20.8 to 24.7°C,  
76 respectively. The mean sunshine hours and mean evaporation (USWB Class-A Open Pan  
77 evaporimeter) ranged from 0.2 to 8.2 hours day<sup>-1</sup> and 2.1 to 5.0 mm per day with an average  
78 of 3.8 hr day<sup>-1</sup> and 3.4 mm per day, respectively.

79 **2.1.2 Initial Soil characteristics**

80

81

82

83

84

85

86

87

Composite soil sample at 0-15 cm depth was collected, processed and analyzed for different physical, chemical properties by following the standard procedures and The soil was sandy clay loam in texture (18.3% clay, 5.5% silt and 76.2% sand), slightly alkaline (7.87) in reaction, non-saline (0.423 dS m<sup>-1</sup>) in nature. The oxidizable organic carbon was medium (6.5 g kg<sup>-1</sup>). The available nitrogen was low (213 kg ha<sup>-1</sup>) and available phosphorus and potassium was in high category (189 and 564 kg ha<sup>-1</sup> respectively) whereas, DTPA extractable zinc was sufficient (1.33 g kg<sup>-1</sup>).

88 **2.2 Treatments and Experimental Design**

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

The experiment was laid out in split plot design with two finger millet varieties as main treatments viz., Vakula and Tirumala released by Agricultural Research Station, Perumallapalli and zinc fertilization at different methods and crop stages as sub treatments viz., T<sub>1</sub>: Control (No fertilizers and manures), T<sub>2</sub>: 60 -30-20 kg N-P-K + FYM @ 4 t ha<sup>-1</sup>, T<sub>3</sub>: T<sub>2</sub> + Soil application of ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> as basal, T<sub>4</sub>: T<sub>2</sub> + Soil application of chelated zinc sulphate @ 5 kg ha<sup>-1</sup>, T<sub>5</sub>:T<sub>2</sub>+Foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence stage, T<sub>6</sub>: T<sub>2</sub> + Foliar application of 0.2% ZnSO<sub>4</sub> at grain filling stage and T<sub>7</sub>:T<sub>2</sub>+Foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence and grain filling stage. The treatments randomized in split plot design with three replications. The recommended dose of 60 kg N, 30 kg P<sub>2</sub>O<sub>5</sub> and 20 kg K<sub>2</sub>O ha<sup>-1</sup> applied through urea, SSP and MOP, respectively. Fertilizer nitrogen was applied in two equal splits as first half dose at the time of transplanting and second half at 30 DAT and the full dose of FYM @ 4 t ha<sup>-1</sup>, phosphorus and potassium applied at the time of transplanting.

103 **2.3 Soil and Plant analysis**

104

105

106

107

108

109

110

111

112

113

Post harvest soil samples were collected from each treatment before after harvesting of both the seasons during 2019 and 2020 at 0 to 15 cm. The samples were air-dried at room temperature, pulverized, sieved through a 2-mm sieve. The available zinc DTPA method (15). Weighed 10 g of soil into a 150 ml conical flask, added 20 ml of DTPA extractant (0.005 M DTPA, 0.01 M CaCl<sub>2</sub> and 0.1 M Triethanol amine (TEA) with pH 7.3) and shake the contents on a horizontal shaker for 2 hrs and Filtered the suspension through whatman No.42 filter paper and zinc content was determined in the extractant using Atomic absorption spectrophotometer (model No: Spectra A varian 220).

114 **2.4 Yield and yield attributes**

115

116

117

118

The grain obtained from the net plot area including the grain of the sampled plants was thoroughly sundried to a safe moisture level of 14 per cent, weighed and expressed in kg ha<sup>-1</sup>. Similarly straw was sun dried to a constant weight and expressed in kg ha<sup>-1</sup>.

119

120

119 **2.5 Estimation of quality parameters**

120

121

122

121 **2.5.1 Grain zinc and iron content (mg kg<sup>-1</sup>)**

123

124

125

126

127

128

Di-acid digestion was carried out using a mixture of HNO<sub>3</sub>:HClO<sub>4</sub> (9:4) by taking one gram of powdered grain sample in 150 ml conical flask and added 10 ml of di-acid mixture and mixed by swirling. The contents were placed on hot plate in a digestion chamber. The contents were further evaporated until the volume was reduced to 3 to 5 ml but not to dryness. The completion of digestion was confirmed by white fumes and kept for cooling. Added double distilled water and filter the contents into a 100 ml volumetric flask by using whatman No. 42

129 filter paper and made upto 100 ml. The filtrate was used for estimating zinc by AAS. Grain  
130 zinc (ppm) = [AAS reading x volume made (100 ml)]/ wt of the plant sample (g).

131

### 132 **2.5.2 Grain protein content (%)**

133 Estimation of total protein content in seeds of finger millet was done as per the  
134 method developed by(16). Weighed 0.5 g grain, grounded with pestle and mortar by adding  
135 10 ml of phosphate buffer. The contents were centrifuged at 3500 rpm for 15 min. The  
136 supernatant was used for protein estimation. Aliquot of 0.2 ml of sample extract was pipette  
137 out in test tube and made up to 1.0 ml volume. A test tube with 1 ml volume of water was  
138 used as a blank. 5 ml of reagent-C was added to all the test tubes including the blank. The  
139 contents were mixed well and allowed to stand for 10 min. added 0.5 ml of reagent-D and  
140 mixed well, incubated for 30 min at room temperature in dark. The colour intensity was read  
141 at 660 nm using spectrophotometer (Model: Genesys 10S UV-VIS). From the standard  
142 curve, concentration of protein in different samples was determined and expressed in  
143 percentage.

144

### 145 **2.6 Statistical Analysis**

146 The experimental data were analyzed statistically by following standard procedure  
147 outlined by (17). Significance was tested by comparing 'F' value at 5 per cent level of  
148 probability. Treatmental differences that were non-significant were denoted as NS and the  
149 data analysed by OPSTAT.

## 150 **3. RESULTS AND DISCUSSION**

### 151 **3.1 Effect of zinc on yield attributes**

152 The yield attributes among the two varieties (main plots) and the interaction effect  
153 were found non-significant. Among the zinc application treatments (sub plots) there is a  
154 significant difference. Two years (2019 & 2020) data was presented. The response was  
155 almost similar among main, sub plots and interactions in two years of experiments and  
156 hence only pooled data are used to highlight the results.

#### 157 **3.1.2 Productive tillers per plant**

158 Number of productive tillers per plant has been tabulated in table 1. From the pooled  
159 data, it was noticed that the productive tiller number per plant was significantly influenced by  
160 the application of zinc. The treatment RDF + FYM +foliar application of 0.2% ZnSO<sub>4</sub> at ear  
161 head emergence and grain filling stages (T<sub>7</sub>) recorded significantly more number of  
162 productive tillers per plant (1.88) which was on par with the treatments RDF + FYM + foliar  
163 application of 0.2% ZnSO<sub>4</sub> at grain filling stage (T<sub>6</sub>) (1.68), RDF + FYM +foliar application of  
164 0.2% ZnSO<sub>4</sub> at ear head emergence stage (T<sub>5</sub>) (1.62) and RDF + FYM +soil application  
165 ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> (T<sub>3</sub>) (1.53) regardless of main plot treatments. The lowest was  
166 expressed in (T<sub>1</sub>) control (1.17). The main plot treatment ranged from 1.43 to 1.61 and there  
167 was no significant difference between the varieties. The interaction effect was noticed as  
168 non-significant and ranged from 1.07 to 1.80. The results are in coincidence with (18) and  
169 (19). Increased number of productive tillers per plant due to optimum supply of zinc which  
170 increases the availability of other nutrients (macro & micro) results in the enhancement of  
171 metabolic activities of plant and finally increased the yield.

#### 172 **3.1.2 No. of finger number per plant**

173 The data pertaining to finger number per plant was presented in table 1. From the pooled  
174 data, The treatment RDF + FYM +foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence  
175 and grain filling stages (T<sub>7</sub>) recorded significantly more number of finger number per plant  
176 (21.58) which was on par with the treatments RDF + FYM + foliar application of 0.2% ZnSO<sub>4</sub>  
177 at grain filling stage (T<sub>6</sub>) (19.40) and RDF + FYM +foliar application of 0.2% ZnSO<sub>4</sub> at ear  
178 head emergence stage (T<sub>5</sub>) (18.17) and the lowest was noticed in (T<sub>1</sub>) control (12.82)  
179 regardless of main plot treatments. There was no significant difference among main plot  
180 treatments and interaction effect. The values ranged between 15.82 to 18.35 and 22.17 to

181 11.00 respectively. Matching results were expressed in the year 2019 and 2020 experiments  
182 conducted by (20) and (21) respectively.

### 183 **3.1.3 Test weight (1000 grains)**

184 The data pertaining to finger number per plant was presented in table 1. From the pooled  
185 data, The treatment RDF + FYM +foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence  
186 and grain filling stages (T<sub>7</sub>) recorded significantly highest test weight (3.05 g) which was on  
187 par with RDF + FYM +foliar application of 0.2% ZnSO<sub>4</sub> at grain filling stage (T<sub>6</sub>) (2.98 g) and  
188 RDF + FYM + foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence stage (T<sub>5</sub>) (2.95 g).  
189 Lowest test weight was registered by absolute control (T<sub>1</sub>) (2.72 g). Among the main plot and  
190 interaction, the effect was non-significant. The values ranged from 2.86 g to 2.94 g and 2.66  
191 g to 3.06 g respectively. The highest test weight (g) in T<sub>7</sub> treatment was due to foliar spray  
192 which helps to rapid absorption of zinc through leaf resulted in better grain filling ability of the  
193 crop. The supremacy of yield components is due to combined application of RDF + FYM +  
194 ZnSO<sub>4</sub> which enhanced the photosynthesis and better translocation of photosynthates from  
195 source to sink. The above results of yield components are in coincidence with the  
196 experimental results conducted by (22) and (23).

### 197 **3.1.4 Grain Yield (kg ha<sup>-1</sup>)**

198 **The data pertaining to grain yield (kg ha<sup>-1</sup>) was presented in table 2.** Grain yield is the  
199 combination of yield attributing characters viz., number of productive tillers per plant, no. of  
200 fingers per plant and test weight (g). From the pooled data, it was revealed that significantly  
201 (**p<0.05**) grain yield (2298.27 kg ha<sup>-1</sup>) was observed in (V<sub>2</sub>) tirumala variety when compared  
202 to vakula variety of finger millet. Among sub plots, the highest grain yield was received by  
203 the treatment applied with RDF + FYM + Foliar application of 0.2% ZnSO<sub>4</sub> at both ear head  
204 emergence and grain filling stages (T<sub>7</sub>) (3150.55 kg ha<sup>-1</sup>). The lowest grain yield of 1452.9 kg  
205 ha<sup>-1</sup> was registered in absolute control (T<sub>1</sub>) compared to other treatments. Regardless of  
206 application of Zinc sulphate, incorporation of FYM + RDF recorded 38% higher when  
207 compared to absolute control. The increase in grain yield in T<sub>7</sub> was 57% compared to T<sub>2</sub>  
208 (RDF + FYM). The interaction effect also shows significant (**p<0.05**) difference and highest  
209 was noticed by Tirumala variety (V<sub>2</sub>) (3048 kg ha<sup>-1</sup>) by the foliar spray of 0.2% ZnSO<sub>4</sub> at both  
210 ear head emergence and grain filling stages (T<sub>7</sub>) *ie.*, (V<sub>2</sub>T<sub>7</sub>) and the lowest (1434 kg ha<sup>-1</sup>)  
211 was noticed by Vakula variety (V<sub>1</sub>) with absolute control (T<sub>1</sub>) *ie.*, V<sub>1</sub>T<sub>1</sub>. Similar results were  
212 reported by (24) and (25)

213

### 214 **3.1.4 Straw yield (kg ha<sup>-1</sup>)**

215 **The data pertaining to straw yield (kg ha<sup>-1</sup>) was presented in table 2.** Application of RDF +  
216 FYM + foliar application of 0.2% ZnSO<sub>4</sub> at ear head emergence and grain filling stages (T<sub>7</sub>)  
217 improved the straw yield (7364.58 kg ha<sup>-1</sup>) of finger millet by 83.2 per cent compared to  
218 absolute control (T<sub>1</sub>) and 51.5 % compared to T<sub>2</sub>. The main plot and interactions were non-  
219 significant and the values are in the range of 5531 to 5423 and 7404 to 3866. kg ha<sup>-1</sup>  
220 respectively. These findings are in matching with (24) and (19). The highest straw yield (kg  
221 ha<sup>-1</sup>) in T<sub>7</sub> treatment was due to increase in cell division and cell elongation. The lowest  
222 straw yield of 3566 kg ha<sup>-1</sup> was expressed by T<sub>1</sub> (absolute control). The grain yield was the  
223 ultimate end product of many yield contributing attributes, physiological and morphological  
224 processes that took place in plants during its life cycle. The Integrated application of organic  
225 and inorganic sources showed beneficial effect on physiological process of plant metabolism  
226 and growth, thereby resulting in higher grain and straw yield.

## 227 **3.2 Quality parameters**

228

### 229 **3.2.1 Protein content in grain (%)**

230 **The data pertaining to protein content (%) in grain was presented in table 3.** There was  
231 significant difference among the zinc application treatments (sub-plots) and the highest  
232 protein content (9.63%) in grain was recorded in the treatment receiving RDF + FYM + foliar  
233 application of 0.2% ZnSO<sub>4</sub> at ear head emergence and grain filling stages (T<sub>7</sub>). Lowest grain

234 yield was registered with absolute control (4.49%). Table. 3. The protein content of two  
235 varieties (main plots) showed non-significant and the values ranged between 7.90 to 7.56%.  
236 The interaction between varieties and zinc application treatments was non-significant (table.  
237 3) on protein content of grain. The values ranged between 9.93% ( $V_2T_7$ ) to 4.54% ( $V_1T_1$ ).  
238 The results were coincided with the results obtained by (22) and (26). The increase in  
239 protein content with zinc application is in agreement with the hypothesis that zinc plays an  
240 important role in protein synthesis.

### 241 **3.3.2 Zinc content in grain ( $\text{mg kg}^{-1}$ )**

242 **The data pertaining to zinc content in grain was presented in table 3.** Among the two  
243 varieties, Tirumala ( $V_2$ ) recorded significantly ( $p < 0.05$ ) highest zinc content ( $28.68 \text{ mg kg}^{-1}$ )  
244 in grain. Among the treatments (zinc application) of sub-plot, ( $T_7$ ) RDF + FYM + foliar  
245 application of 0.2%  $\text{ZnSO}_4$  at ear head emergence and grain filling stages recorded  
246 significantly ( $p < 0.05$ ) highest zinc content ( $39.06 \text{ mg kg}^{-1}$ ) in grain. The lowest was  
247 registered by absolute control ( $T_1$ ) ( $12.02 \text{ mg kg}^{-1}$ ). The interaction effect shows non-  
248 significant results and values ranged between 11.17 to  $40.82 \text{ mg kg}^{-1}$ . The results are in  
249 coincidence with the experiments conducted during the year 2019 and 2020 by (27) and (28)  
250 respectively. Enhancement of zinc content in grain of finger millet by 69.5% compared to  
251 RDF + FYM ( $T_2$ ) was due to foliar application of 0.2%  $\text{ZnSO}_4$  at both ear head emergence  
252 and grain filling stages. Whereas, the increase of protein content of  $T_7$  by (69.5%) compared  
253 to  $T_2$

### 254 **3.3.3 Iron content in grain ( $\text{mg kg}^{-1}$ )**

255 **The data pertaining to iron content in grain was presented in table 3.** The iron content ( $\text{mg}$   
256  $\text{kg}^{-1}$ ) in grain was significantly influenced by sub plots (zinc application) and main plot  
257 (varieties) and interaction between varieties and zinc application showed non-significant.  
258 Highest iron content ( $151.81 \text{ mg kg}^{-1}$ ) was registered significantly with Tirumala variety ( $V_2$ ).  
259 Significantly highest iron content ( $178.92 \text{ mg kg}^{-1}$ ) in grain was recorded by  $T_7$  (RDF + FYM  
260 + foliar application of 0.2%  $\text{ZnSO}_4$  at ear head emergence and grain filling stages) which  
261 was on par with  $T_6$  ( $168.91 \text{ mg kg}^{-1}$ ) (RDF + FYM + foliar application of 0.2%  $\text{ZnSO}_4$  at grain  
262 filling stage). Both the treatments were significantly superior over other treatments which  
263 were tried in this experiment. The results are similar with the field experiments conducted by  
264 (29) and (30). The Lowest iron content ( $114.28 \text{ mg kg}^{-1}$ ) in grain of finger millet was recorded  
265 with absolute control ( $T_1$ ). The interaction effect ranged from 182.07 to  $113.80 \text{ mg kg}^{-1}$  in  
266 grains of finger millet. The iron content was increased by (43.2%) compared to RDF + FYM  
267 ( $T_2$ ) whereas the increase in iron content by 55.2% compared to absolute control ( $T_1$ ).  
268 The foliar application of  $\text{ZnSO}_4$  at both ear head emergence and grain filling stages  
269 enhanced all grain quality parameters.

270  
271  
272

**Table 1. Yield and yield attributes as influenced by zinc application of finger millet varieties**

| Treatments  | No. of productive tillers per plant |       |        | Finger number per plant |       |        | Test weight (1000 grain weight) |       |        |
|---|-------------------------------------|-------|--------|-------------------------|-------|--------|---------------------------------|-------|--------|
|   | 2019                                | 2020  | Pooled | 2019                    | 2020  | Pooled | 2019                            | 2020  | Pooled |
| <b>Varieties (V)</b>  |                                     |       |        |                         |       |        |                                 |       |        |
| (V <sub>1</sub> ) : Vakula  | 1.97                                | 0.86  | 1.43   | 16.81                   | 15.17 | 15.98  | 2.95                            | 2.76  | 2.86   |
| (V <sub>2</sub> ) : Tirumala  | 2.26                                | 0.95  | 1.61   | 18.80                   | 17.89 | 18.35  | 2.98                            | 2.90  | 2.94   |
| S.E (m)   | 0.125                               | 0.044 | 0.062  | 1.061                   | 1.015 | 1.048  | 0.040                           | 0.014 | 0.025  |
| C.D (p=0.05)  | NS                                  | NS    | NS     | NS                      | NS    | NS     | NS                              | 0.095 | NS     |
| <b>Method and stage of Zn application</b>   |                                     |       |        |                         |       |        |                                 |       |        |
| Control (no fertilizers and manures)  | 1.66                                | 0.68  | 1.17   | 13.08                   | 12.55 | 12.82  | 2.76                            | 2.68  | 2.72   |
| RDF (60 -30-20 kg N-P-K & FYM @ 4 t/ha)   | 1.85                                | 0.75  | 1.30   | 14.98                   | 13.45 | 14.22  | 2.93                            | 2.77  | 2.86   |
| RDF+ Soil application of ZnSO <sub>4</sub> @ 25 kg/ha as basal                                | 2.18                                | 0.87  | 1.53   | 17.93                   | 16.73 | 17.33  | 2.93                            | 2.85  | 2.89   |
| RDF+ Soil application of chelated Zinc @ 5 kg/ha  | 2.07                                | 0.83  | 1.47   | 17.33                   | 15.95 | 16.63  | 2.90                            | 2.80  | 2.85   |
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at ear head emergence stage                 | 2.22                                | 0.97  | 1.62   | 18.90                   | 17.45 | 18.17  | 3.02                            | 2.87  | 2.95   |
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at grain filling stage                      | 2.33                                | 0.98  | 1.68   | 19.98                   | 18.78 | 19.40  | 3.06                            | 2.89  | 2.98   |
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at ear head emergence & grain filling stage | 2.51                                | 1.25  | 1.88   | 22.40                   | 22.78 | 21.58  | 3.17                            | 2.95  | 3.05   |





**Table 2. Grain yield and straw yield as influenced by zinc application in finger millet varieties**

| Treatments  | Grain yield (kg ha <sup>-1</sup> ) |       |        | Straw yield (kg ha <sup>-1</sup> ) |       |        |
|---|------------------------------------|-------|--------|------------------------------------|-------|--------|
|   | 2019                               | 2020  | Pooled | 2019                               | 2020  | Pooled |
| <b>Varieties (V)</b>  |                                    |       |        |                                    |       |        |
| (V <sub>1</sub> ) : Vakula  | 2481                               | 1980  | 2230   | 5922                               | 4923  | 5423   |
| (V <sub>2</sub> ) : Tirumala  | 2585                               | 2010  | 2298   | 6195                               | 4867  | 5531   |
| S.E (m)   | 6.1                                | 6.7   | 6.03   | 142.7                              | 188.2 | 146.4  |
| C.D (p=0.05)  | 40.0                               | NS    | 40.2   | NS                                 | NS    | NS     |
| <b>Method and stage of Zn application</b>   |                                    |       |        |                                    |       |        |
| Control (no fertilizers and manures)  | 1666                               | 1239  | 1452   | 4641                               | 3395  | 4018   |
| RDF (60 -30-20 kg N-P-K & FYM @ 4 t/ha)   | 2183                               | 1823  | 2003   | 5262                               | 4454  | 4858   |
| RDF+ Soil application of ZnSO <sub>4</sub> @ 25 kg/ha as basal                                | 2413                               | 1990  | 2201   | 5516                               | 4662  | 5089   |
| RDF+ Soil application of chelated Zinc sulphate @ 5 kg/ha                                     | 2353                               | 1933  | 2143   | 5366                               | 4591  | 4979   |
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at ear head emergence stage                 | 2521                               | 2073  | 2297   | 6691                               | 4866  | 5779   |
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at grain filling stage                      | 3036                               | 2169  | 2603   | 6908                               | 5595  | 6252   |
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at ear head emergence & grain filling stage | 3560                               | 2740  | 3150   | 8025                               | 6704  | 7364   |
| S.E (m)   | 76.8                               | 25.7  | 40.5   | 335.7                              | 236.8 | 190.8  |
| C.D (p=0.05)  | 22.5                               | 75.4  | 119.0  | 985.8                              | 695.4 | 560.2  |
| Interaction (main x sub)  | NS                                 | 111.8 | 171.1  | NS                                 | NS    | NS     |


1 **Table 3. Grain quality parameters of finger millet varieties as influenced by zinc application**

| Treatments  | Protein (%) |       |        | Zinc (mg kg <sup>-1</sup> ) |       |        | Iron (mg kg <sup>-1</sup> ) |       |        | Calcium (%) |      |        |
|---|-------------|-------|--------|-----------------------------|-------|--------|-----------------------------|-------|--------|-------------|------|--------|
|   | 2019        | 2020  | Pooled | 2019                        | 2020  | Pooled | 2019                        | 2020  | Pooled | 2019        | 2020 | Pooled |
| <b>Varieties (V)</b>  |             |       |        |                             |       |        |                             |       |        |             |      |        |
| (V <sub>1</sub> ) : Vakula  | 7.70        | 7.41  | 7.56   | 25.10                       | 24.06 | 24.58  | 149.38                      | 147.5 | 148.4  | 0.27        | 0.68 | 0.48   |
| (V <sub>2</sub> ) : Tirumala  | 8.10        | 7.69  | 7.90   | 29.39                       | 27.97 | 28.68  | 152.93                      | 150.6 | 151.8  | 0.29        | 0.74 | 0.51   |
| S.E (m)   | 0.09        | 0.053 | 0.07   | 0.96                        | 0.72  | 0.25   | 2.81                        | 4.6   | 1.92   | 0.007       | 0.17 | 0.09   |
| C.D (p=0.05)  | NS          | NS    | NS     | NS                          | NS    | 1.682  | NS                          | NS    | NS     | NS          | NS   | NS     |
| <b>Method and stage of Zn application</b>                                     |             |       |        |                             |       |        |                             |       |        |             |      |        |
| Control (no fertilizers and manures)  | 4.61        | 4.36  | 4.49   | 12.69                       | 11.36 | 12.02  | 116.2                       | 114.2 | 115.2  | 0.30        | 0.52 | 0.41   |
| RDF (60 -30-20 kg N-P-K & FYM @ 4 t/ha)                                       | 6.89        | 6.78  | 6.84   | 23.66                       | 22.44 | 23.05  | 123.0                       | 126.7 | 124.8  | 0.27        | 0.57 | 0.42   |
| RDF+ Soil application of ZnSO <sub>4</sub> @ 25 kg/ha as basal                | 8.62        | 8.35  | 8.48   | 27.47                       | 25.88 | 26.68  | 156.3                       | 154.2 | 155.3  | 0.28        | 0.97 | 0.63   |
| RDF+ Soil application of chelated Zinc sulphate @ 5 kg/ha                     | 8.34        | 8.07  | 8.21   | 25.13                       | 24.29 | 24.71  | 149.1                       | 144.2 | 146.6  | 0.30        | 0.87 | 0.59   |
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at ear head emergence stage | 8.16        | 7.64  | 7.90   | 28.36                       | 27.36 | 27.86  | 162.7                       | 159.3 | 161.0  | 0.27        | 0.40 | 0.33   |
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at grain filling stage      | 8.69        | 8.37  | 8.53   | 33.54                       | 32.55 | 33.04  | 169.9                       | 167.9 | 168.9  | 0.26        | 0.78 | 0.52   |

|   |       |       |       |       |       |       |       |       |       |       |       |       |
|---|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| RDF+ foliar application of 0.2% ZnSO <sub>4</sub> at ear head emergence & grain filling stage | 9.97  | 9.30  | 9.63  | 39.87 | 38.25 | 39.06 | 180.8 | 177.0 | 178.9 | 0.29  | 0.85  | 0.57  |
| S.E (m)   | 0.334 | 0.244 | 0.277 | 1.617 | 1.459 | 1.084 | 7.859 | 7.207 | 5.567 | 0.020 | 0.210 | 0.107 |
| C.D (p=0.05)  | 0.981 | 0.715 | 0.813 | 4.747 | 4.284 | 3.184 | 23.0  | 21.16 | 16.34 | NS    | NS    | NS    |
| Interaction(main x sub)   | NS    | NS    | NS    | NS    | NS    | NS    | NS    | NS    | NS    | 0.089 | NS    | NS    |

2  
3  
4

**Graph 1. Influence of zinc application on grain zinc content in finger millet**



5

#### 6 4. CONCLUSION

7  
8 The results of the present experiment confess that significantly yield attributes, yield  
9 and quality parameters of finger millet were recorded highest with the application of RDF+  
10 foliar spray of 0.2% ZnSO<sub>4</sub> at both ear head emergence and grain filling stages. It might be  
11 acknowledged that combination of inorganic and organic nutrient sources and addition of  
12 external application of zinc through foliar spray proved to be superior over other treatments.  
13 So, we can informed that the application of zinc sulphate at both stages enhanced the yield,  
14 yield parameters and quality parameters compared to other treatments of tirumala variety  
15 under rainfed alfisols of southern zone of Andhra Pradesh.

#### 16 17 ACKNOWLEDGEMENTS

18 All authors of this manuscript are thankful Acharya NG Ranga Agricultural University for  
19 providing financial assistance to carryout research work

#### 20 21 COMPETING INTERESTS

22 Authors have declared that no competing interests exist.

#### 23 24 REFERENCES

- 25  
26 1. Chethan S. Nagappa Malleshi G. Finger Millet Polyphenols: Characterization and  
27 their Nutraceutical Potential. American Journal of Food Technology. 2007;2(7): 2167  
28 – 2179.  
29 2. Ministry of Agriculture, Government of India Indiatat. 2017-18  
30 [www.indiaagrstat.2017-18](http://www.indiaagrstat.2017-18).  
31 3. Saravanpandian P, Subramanian S, Paramasivam P, Kumaraswamy K. Organic  
32 farming in sustaining soil health. Agricultural Reviews. 2005; 26(2): 141 – 147.  
33 4. Shanmugan A. Stanly Joseph Michaelraj PA. Study on millets cultivation in Karur  
34 district of Tamilnadu. International Journal of Management Research & Review.  
35 2013;3(1):2167-2179.  
36 5. Erenso D, Asfaw A, Taye T, Tesfaye T. Genetic resources, breeding and production  
37 of millets in Ethiopia. Proceedings of an International Conference, 19-21 September,  
38 Bern, Switzerland. 2009;43-56.  
39 6. Auld DS. Zinc coordination sphere in biochemical zinc sites, biometals Australian  
40 soils. Australian Journal of Soil Research. 2001;30:45-53  
41 7. Fischer WC, Black RE. Zinc and the risk for infectious disease. Annu Rev Nutr.  
42 2004;24: 255–75.  
43 8. De Valença AW, Bake A, Brouwer ID, Giller KE. Agronomic biofortification of crops  
44 to fight hidden hunger in sub-Saharan Africa. Global Food Secur. 2017;12:8–14.  
45 9. Liu J, Yang M, Li H, Li D, Shi X, Zhang Y. Genetic processes of iron and zinc  
46 accumulation in edible portion of crops and their agro-biofortification: a review. Am J  
47 Agric For. 2017;5(3):65–72.  
48 10. Singh MV. Micro and Secondary Nutrients and Pollutant Elements Research in  
49 India. Progress Report, IISS, Bhopal, 2006;1-110.  
50 11. Ramegowda Y, Venkategowda R, Govind G, Hagadur GJ, Udayakumar M, Shankar  
51 AG. Effect of Zn application on its uptake, distribution and concentration of Fe and  
52 Cu in finger millet (*Eleusine coracana* (L.) Gaertn). Journal of Plant Nutrition. 2016;  
53 39(4): 569-580  
54 12. Shivay YS, Kumar D, Prasad R, Ahlawat IPS. Relative yield and zinc uptake by rice  
55 from zinc sulphate and zinc oxide coatings onto urea. Nutr cyc agroecosyst.  
56 2008;80:181-188.

- 57 13. Yang XW, Tian XH, Gale WJ, Cao YM, Lu XC, Zhao AQ. Effect of soil and foliar  
58 application on zinc concentration and bioavailability in wheat grain grown on  
59 potentially zinc deficient soils. *Cereal Res Commun.* 2011;39: 553-543.
- 60 14. Hossain MA, Jahiruddin M, Islam MR, Mian MH. The requirement of zinc for  
61 improvement of crop yield and mineral nutrition in maize-mungbean-rice system.  
62 *Plant soil.* 2008;306: 13-22.
- 63 15. Lindsay WL, Norvell WA. Developments of DTPA soil test for zinc, iron, manganese  
64 and copper. *Soil Science Society of American Journal.* 1978;42: 421-428
- 65 16. Lowry OH, Rosenbrough NJ, Farr AL, Randall RJ. Protein measurement with the  
66 folin phenol reagent. *Journal of Biochemistry.* 1951;193: 265-375.
- 67 17. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*, 1985.ICAR,  
68 New Delhi
- 69 18. Patel NI, Patel BR, Patel CK and Patel FB. Effect of foliar spray of nutrients at  
70 different growth stages on pearl millet under Dryland condition. *International Journal*  
71 *of Chemical Studies.* 2019;7(5): 05-09.
- 72 19. Logesh Kumar S, Sankaran, VM, Hemalatha M and Suresh S. Effect of different  
73 level of fertilizers and foliar nutrition on yield attributes and yield of Pearl millet under  
74 rainfed condition. *International Journal of Chemical Studies.* 2020; 8(6): 979-982.
- 75 20. Kamal G. Comparative response of finger millet [*Eleusine coracana* (L.) Gaertn] and  
76 pearl millet [*Pennisetum glaucum* (L.) R.Br. emend Stuntz] to applied nutrients under  
77 semi arid conditions of Rajasthan. M.Sc(Ag.)Thesis submitted to Sri Karan Narendra  
78 Agriculture University, Jobner, India. 2019.
- 79 21. Vijaya kumar M, Sivakumar M and Tamilselvan N. Effect of Zinc and Iron application  
80 on yield attributes, available nutrients status and nutrient uptake of finger millet  
81 under rainfed condition. *International Journal of Current Microbiology and Applied*  
82 *Sciences.* 2020;9(5): 3237 – 3246.
- 83 22. Akshay G. Effect of zinc biofortification on soil properties, yield, nutrient uptake and  
84 quality of *rabi* sorghum. M.Sc(Ag.)Thesis submitted to Vasantrya Naik Marathwada  
85 Krishi Vidyapeeth, Parbhani, India. 2019.
- 86 23. Prashantha GM, Prakash SS, Umesh S, Chikkaramappa T, Subbarayappa CT and  
87 Ramanamurthy V. Direct and residual effect of zinc and boron on yield and yield  
88 attributes of finger millet – groundnut cropping system. *International Journal of Pure*  
89 *and Applied Bioscience.* 2019;7(1): 124 – 134.
- 90 24. Ashoka P, Rajakumar GR and Priya P. Response of soil and foliar application of  
91 nutrients on yield and nutrient uptake of foxtail millet (*Setaria italica* L.). *International*  
92 *Journal of Chemical Studies.* 2020; 8(4): 2282-2285.
- 93 25. Chowdary KA and Patra BC. Effect of Micronutrient application with different  
94 sources of NPK on growth and yield of finger millet crop in red laterite zone. *Journal*  
95 *of Agricultural Science and Technology.* 2019;9: 403-416.
- 96 26. Sharad KM, Suresh K, Madhavi A and Shanti M. Agronomic biofortification and  
97 nutrient uptake of sorghum (*Sorghum bicolor* L.) as influenced by fertilization  
98 strategies. *International Journal of Chemical Studies.* 2020; 8(3): 2107-2110.
- 99 27. Sharanappa HS, Latha BK, Desai B, Koppalkar G and Ravi MV. Effect of Agronomic  
100 Biofortification with Zinc and Iron on Yield and Quality of Pearl millet [*Pennisetum*  
101 *glaucum* (L.)] Genotypes. *International Journal of Current Microbiology and Applied*  
102 *Sciences.* 2019; 8(9): 1312-1321.
- 103 28. Ajay Kumar E, Surekha K, Bhanu K, Rekha and Harish Kumar Sharma S. Effect of  
104 Various Sources of Zinc and Iron on Grain Yield, Nutrient Uptake and Quality  
105 Parameters of Finger Millet (*Eleusine coracana* L.). *International Research Journal*  
106 *of Pure & Applied Chemistry.* 2020; 21(2): 46-55.
- 107 29. Amar KJ, Shrivastava S and Vinay A. Response of organic manure, zinc and iron on  
108 soil properties, yield and nutrient uptake by Pearl millet crop grown in inceptisol.  
109 *International Journal of Pure and Applied Bioscience.* 2018; 6 (1): 426-435.

- 110 30. Sravani K. Zinc fortification for grain quality and yield of finger millet (*Eleusine*  
111 *Coracana* (L.) Gaertn). M.Sc(Ag.)Thesis submitted to Acharya N G Ranga  
112 Agricultural University, Hyderabad, India. 2018.  
113  
114  
115