

## Original Research Article

# Qualitative Analysis of the Phytochemical Contents of Different Anatomical Parts of Ripe *Solanumaethiopicum* Linn Fruits

### ABSTRACT

**AIMS:** To determine qualitatively the amount of alkaloids, saponin, tannin, volatile oil, phenol and flavonoids in the different anatomical parts of ripe *Solanumaethiopicum* Linn fruits.

**METHODOLOGY:** *Solanumaethiopicum* Linn fruits were purchased from Sabo market, Ogbomoso, Nigeria. The fruits were separated with a razor into four (4) anatomical parts (the epicarp, mesocarp, endocarp, and seed) after washing with distilled water. 1gram of each anatomical parts of the fruit were soaked in 20mls of distilled water, 1% volume per volume (v/v), 2% v/v, 3%v/v, 4%v/v and 5%v/v of ethanol, ethylacetate and methanol separately. The mixtures were left for 24 hours after which they were sieved to obtain the extracts. The presence of alkaloid, tannin, saponin, volatile oil, phenol and flavonoid were determined qualitatively in the extracts at selected concentrations.

**RESULT:** The results obtained showed that alkaloids are more concentrated in the mesocarp of *Solanumaethiopicum* Linn; volatile (essential) oil is more pronounced in the mesocarp and endocarp of *Solanumaethiopicum* Linn; saponin is confirmed in all anatomical parts of the fruit at reasonable quality except in the seed; tannin is found to be moderately present only in the aqueous extract of mesocarp of *Solanumaethiopicum* Linn fruit while the presence of phenol and flavonoids were confirmed in trace amount at few tested concentrations of the extracts.

**CONCLUSION:** Taken together, the presence of a variety of phytochemicals in the different anatomical parts of *Solanumaethiopicum* Linn fruit indicate that the fruit might be pharmacologically active against a number of diseases. However, this should be subjected to subsequent researches

Keywords: *Solanumaethiopicum* Linn, garden egg, phytochemical content, epicarp, mesocarp, endocarp, seed.

## 1.0 INTRODUCTION

*Solanumaethiopicum* Linn. popularly known as garden egg or African eggplants are highly consumed for dietary and medicinal purposes in most African countries. It is known as dauta, afufa or añara, and igbagbain Hausa, Igbo and Yoruba Languages respectively which are the major languages in Nigeria [1].

### Scientific classification

Kingdom: Plantae

Order: Solanales

Family: Solanaceae

Genus: Solanum

Species: *S. aethiopicum*

Authority: Linneaus

Botanical name: *Solanumaethiopicum* LinnLinn

### Botanical description

Different species of the plant have been shown to demonstrate a level of variation in certain features which includes diameter of petal, petiole length, leaf blade width, plant branching, fruit shape and colour [1]. Garden egg fruit is round in shape with flattened out top and bottom and grooved portions which are averagely 5-6cm long and 6-7cm wide. It usually has a tiny seed and straight or arc stalk [2].



FIG.1 *Solanumaethiopicum Linn* fruit

Garden egg fruit can be categorized anatomically into three distinct layers namely the epicarp (also known as exocarp), which is the outermost layer; the mesocarp, which is the middle layer; and the endocarp (inner layer surrounding the seeds).




FIG.2 Anatomical parts of *Solanumaethiopicum Linn* fruit

## Traditional and Pharmacological activities

Traditionally, *Solanumaethiopicum Linn* fruits are used in the treatment of a number of diseases which include catarrh, asthma, allergic rhinitis, skin infections, gastro-esophageal reflux disease, constipation, obesity, **dyspepsia**, rheumatic disease and swollen joint pains, amidst others [3, 4]. Its pharmacological activities include antidiabetic activity, anti-hypertensive activity [5], anti-inflammatory activity, antiulcer activity [6], antioxidant [7] and anti-obesity effect [8]. Other reported pharmacological activities are analgesic, anti-asthmatic, anti-glaucoma, hypoglycemic, hypolipidemic and weight reduction effects on animals and human [9, 10]. The pharmacological activities of *Solanumaethiopicum Linn* are said to be connected to its phytochemical constituents [11].

## Phytochemical profile

Phytochemicals are naturally occurring plant's bio-active constituents which provide health benefits for mankind [12]. Plants produce these chemicals primarily to protect themselves and secondarily humans against diseases [13]. They also contribute to the plant's features such as colour, aroma and taste [14]. Phytochemicals can be found in different parts of the plants, such as the leaves, flowers, roots, stems, seeds and fruits however their concentration varies from plant to plant depending on variety, growth conditions, season and so on [13]. Phytoconstituents reported to be present in ***Solanumaethiopicum Linn*** include alkaloids, flavonoids, phytosterols, saponins, ascorbic acid, cardiac glycosides, tanins and terpenols amidst others [15; 16].

Up to date, there is limited information as regards the phytochemical **contents** of the different anatomical parts of *Solanumaethiopicum Linn* fruit hence the need for this research

## 2.0 MATERIALS AND METHODS

### 2.1 Chemical and reagents

Mercury chloride, potassium iodide, sodium hydroxide, hydrochloric acid (dilute and concentrated), iron (iii) chloride, ethanol, ethylacetate, methanol.

### 2.2 Plant material and extraction

*Solanumaethiopicum Linn* fruits were purchased from Sabo market, Ogbomoso, along Ilorin Express road, Oyo state, Nigeria. The fruits were selected and thoroughly washed with water to remove unwanted particles after which it was separated with a razor into four (4) anatomical parts (epicarp, mesocarp, endocarp and seed) which were used for the experiment. 1gram of each anatomical parts of the fruit were soaked in 20mls of distilled water, 1% volume per volume (v/v), 2% v/v, 3%v/v, 4%v/v and 5%v/v of ethanol, ethylacetate and methanol separately. The mixture was left for 24 hours after which they were sieved to obtain the extracts.

## **2.3 Methodology**

### **2.3.1 Test for Alkaloid**

Two drops of Mayer's reagent together with two drops of 2% v/v of Hydrochloric (HCl) acid was added to 1ml of plant extracts. The cream colouration of the mixture indicates the presence of Alkaloid [17].

### **2.3.2 Test for Volatile oil**

0.2mls of 1% v/v of sodium hydroxide was added to 1ml of plant extracts. Formation of white precipitate indicates the presence of Volatile oil [18].

### **2.3.3 Test for Saponin**

1ml of distilled water was added to 1mls of plant extracts. Foaming after vigorous shaking for about 20minutes indicates the presence of Saponin[19].

### **2.3.4 Test for Phenol and tannin**

Two drops of  $\text{FeCl}_3$  was added to a mixture of 1ml of distilled and 1ml of plant extracts. The formation of a green precipitate indicates the presence of tannin and bluish green or dark green indicates the presence of phenol [20].

### **2.3.5 Test for Flavonoid**

3 drops of 20% NaOH was added to the mixture of 1ml of distilled and 1ml of plant extracts. The intense yellow colouration of the mixture indicates the presence of Flavonoid. The disappearance of the yellow colour after the addition of 1mls of dilute HCl acid validates the presence of Flavonoid [21].

### 3.0 RESULTS AND DISCUSSION

The phytochemical screening of *Solanumaethiopicum* Linn (seed) as shown in table 1 shows that volatile oil, phenol, tannin and flavonoid were absent at all tested concentrations of the extracts (i.e. aqueous, ethanol, ethyl acetate, methanol extracts). Alkaloid was present only in 2% methanol extracts of *Solanumaethiopicum* Linn (seed). Saponin was present in 2% ethanol, 3% ethyl acetate and 3% methanol extracts.

**Table 1: Phytochemical test for *Solanumaethiopicum* Linn (seed)**

EXTRACTS	<i>Solanumaethiopicum</i> Linn (Seed)					
	Alkaloid	Volatile oil	Saponin	Phenol	Tanin	Flavonoid
Aqueous	-	-	-	-	-	-
1%Ethanol	-	-	-	-	-	-
2%Ethanol	-	-	+	-	-	-
3%Ethanol	-	-	-	-	-	-
4%Ethanol	-	-	-	-	-	-
5%Ethanol						
1%Ethyl-acetate	-	-	-	-	-	-
2%Ethyl-acetate	-	-	-	-	-	-
3% Ethyl-acetate	-	-	+	-	-	-
4%Ethyl-acetate	-	-	-	-	-	-
5%Ethyl-acetate	-	-	-	-	-	-
1%Methanol	-	-	-	-	-	-
2%Methanol	+	-	-	-	-	-
3%Methanol	-	-	+	-	-	-
4%Methanol	-	-	-	-	-	-
5%Methanol	-	-	-	-	-	-

**Table 2: Phytochemical test for *Solanumaethiopicum* Linn (mesocarp)**

EXTRACTS	<i>Solanumaethiopicum</i> Linn (Mesocarp)					
	Alkaloid	Volatile oil	Saponin	Phenol	Tanin	Flavonoid
Aqueous	++	++	++	-	++	-
1%Ethanol	++	++	-	-	-	-
2%Ethanol	-	+	+	-	-	-
3%Ethanol	-	-	++	-	-	-
4%Ethanol	+++	-	-	-	-	-
5%Ethanol	+	-	+	-	-	-
1%Ethyl-acetate	-	-	-	-	-	-
2%Ethyl-acetate	++	-	-	-	-	+
3% Ethyl-acetate	+	-	+	-	-	-
4%Ethyl-acetate	++	-	+	-	-	-
5%Ethyl-acetate	++	-	-	-	-	-
1%Methanol	++	-	-	-	-	-
2%Methanol	-	-	-	-	-	-
3%Methanol	-	-	-	-	-	-
4%Methanol	-	-	-	-	-	-
5%Methanol	+	-	-	-	-	-

According to

table 2, alkaloid was present in 5% ethanol extract, 3% ethylacetate extract and 5% methanol extract; moderately present in aqueous extract, 1% ethanol extract, 2%, 4% and 5% ethylacetate extracts, 1% methanol extract and highly present in 4% ethanol extract of *Solanumaethiopicum* Linn (mesocarp). Volatile oil was found to be present only in 2% ethanol extract and moderately present in aqueous and

1%ethanol extracts of *Solanumaethiopicum* Linn (mesocarp). Phenol was absent at all tested concentrations of the extracts. Saponin was present in 2% and 5% ethanol extracts, 3% and 4% ethylacetate extracts and moderately present in aqueous and 3% ethanol extracts. Tannin was moderately present in aqueous extract while flavonoid was only present in 2%ethylacetate extract.

**Table 3: Phytochemical test for *Solanumaethiopicum* Linn(epicarp)**

EXTRACTS	<i>Solanumaethiopicum</i> Linn (Epicarp)					
	Alkaloid	Volatile oil	Saponin	Phenol	Tanin	Flavonoid
Aqueous	-	-	-	-	-	-
1%Ethanol	-	-	+	-	-	-
2%Ethanol	-	-	-	-	-	-
3%Ethanol	-	-	+	-	-	-
4%Ethanol	-	-	-	-	-	-
5%Ethanol	-	-	-	-	-	-
1%Ethyl-acetate	-	-	++	-	-	-
2%Ethyl-acetate	-	-	+	-	-	-
3% Ethyl-acetate	-	-	+	-	-	-
4%Ethyl-acetate	-	-	++	-	-	-
5%Ethyl-acetate	-	-	+	-	-	-
1%Methanol	-	-	-	-	-	-
2%Methanol	-	-	+	-	-	+
3%Methanol	-	-	+	-	-	-
4%Methanol	-	-	-	-	-	-
5%Methanol	-	-	-	-	-	-

As shown in table 3, alkaloid, volatile oil, phenol and tanin were absent at all concentrations of the selected extracts of *Solanumaethiopicum* Linn (Epicarp).Flavonoid is found to be present only in 2%



methanol extract. Saponin was present in 1% and 3% ethanol extracts, 2%, 3% and 5% ethylacetate extracts, 2% and 3% methanol extracts and moderately present in 1% and 4% ethylacetate extracts.

**Table 4: Phytochemical test for *Solanumaethiopicum* Linn (Endocarp)**

EXTRACTS	<i>Solanumaethiopicum</i> Linn (Endocarp)					
	Alkaloid	Volatile oil	Saponin	Phenol	Tanin	Flavonoid
Aqueous	-	-	+	+	-	-
1%Ethanol	+	+	+	-	-	-
2%Ethanol	-	-	++	-	-	-
3%Ethanol	-	++	++	-	-	-
4%Ethanol	-	-	+	-	-	-
5%Ethanol	-	-	++	-	-	-
1%Ethyl-acetate	+	+	+++	-	-	-
2%Ethyl-acetate	-	-	+	-	-	-
3% Ethyl-acetate	-	-	++	-	-	-
4%Ethyl-acetate	-	-	++	-	-	-
5%Ethyl-acetate	-	-	+	-	+	-
1%Methanol	-	-	-	-	-	-
2%Methanol	-	-	-	-	-	-
3%Methanol	+	+	-	+	-	-
4%Methanol	-	-	+	-	-	-
5%Methanol	-	-	++	-	-	-

Table 4 above shows that alkaloid was present in 1% ethanol extract, 1% ethylacetate extract and 3% methanol extract while tannin is only present in 5% ethylacetate extract. Saponin was present in aqueous extract, 1% and 4% ethanol extracts, 2% and 5% ethylacetate extracts and 4% methanol extract; moderately present in 2%, 3% and 5% ethanol extracts, 3% and 4% ethylacetate extracts and 5%

methanol extract; highly present in 1% ethylacetate extract. Volatile oil was found to be present in 1% ethanol extract, 1% ethylacetate extract, 3% methanol extract and moderately present in 3% ethanol while flavonoid was completely absent at all tested concentrations of the selected extracts of *Solanumaethiopicum* Linn (Endocarp).

+ = Present in trace amount

++ = Present in moderate amount

+++ = Present in high concentration

– = Absent

Phytochemical screening is a means of determining qualitatively or quantitatively the amount of substance of therapeutic and nutritional benefit in plants. Plants that are rich in certain phytoconstituents have always been of pharmacological importance. In this study, the phytochemical contents of the different anatomical parts of *Solanumaethiopicum* Linn fruit were determined qualitatively to know which part is rich in the tested phytochemicals at selected extracts.

The presence of alkaloids, volatile oil, saponin, phenol, flavonoid and tannin were confirmed in the different anatomical part of *Solanumaethiopicum* Linn fruit albeit at different concentration. This observation is in line with the studies of various scientists who have confirmed the presence of the phytochemicals in the *Solanumaethiopicum* Linn fruit in previous times although their analysis focused on the whole fruit [11, 15, 16, 22, 23].

Comparing the tables above, it is observed that alkaloids are more concentrated in the mesocarp of *Solanumaethiopicum* Linn. Alkaloids are said to be pharmacologically active against certain diseases such as malaria, body pain, hypertension and cell metastasis amidst others [24, 25]. Hence, this research suggests that the mesocarp of *Solanumaethiopicum* Linn fruit might be useful in the treatment of these diseases.

As shown in the tables above, the presence of volatile (essential) oil is more pronounced in the mesocarp and endocarp of *Solanumaethiopicum* Linn fruit compared to other anatomical parts, however at few tested concentrations. Since essential oils and their components possess antibacterial, antifungal,

antiviral, insecticidal, and antioxidant properties [26, 27], these parts of the plant might possess antimicrobial and free radical scavenging properties. Moreover, plant volatile oils have been said to possess other therapeutic activities which include antibiotic, analgesic, anti-inflammatory, antiseptic, astringent, diuretic, laxative, sedative, vasodilative and vasoconstrictive effects amidst others [28, 29].

A comparison of the above tables shows the presence of saponin in all anatomical parts of *Solanumaethiopicum Linn* fruit at reasonable quality except in the seed where it is only observed in trace amount. It is more pronounced in the endocarp of *Solanumaethiopicum Linn*. The biological effect exerted by saponin includes antimicrobial, anti-tumour, anti-insect, hepatoprotective, haemolytic and anti-inflammatory activities. They cause a decrease in blood cholesterol level and may be used as an adjuvant in vaccines [30].

Tannin is found to be moderately present in the aqueous extract of mesocarp of *Solanumaethiopicum Linn* fruit. Research has shown that Japanese and Chinese uses tannin-containing plant extracts as diuretics against stomach and duodenal tumours [31], and as an anti-inflammatory, antiseptic and antioxidant [32]. Intake of tannins was affirmed by Serrano and colleagues to decrease the frequency of chronic disease [33]. Dietary consumption of tannin also helps in treating hemochromatosis [34].

The presence of phenol and flavonoids are only confirmed in trace amount at a few tested concentrations of the extracts.

#### **4.0 CONCLUSION**

Taken together, the presence of a variety of phytochemicals in the different anatomical parts of *Solanumaethiopicum* fruit indicate that the fruit might be pharmacologically active against a number of diseases. However, this should be subjected to subsequent researches.

#### **Ms\_JOCAMR\_56868 REFERENCES**

1. Childer NF, Margoles MS. An Apparent Relation of Mightshades Solanaceae to Arthritis. Journal of Neurological and Orthopedic Medical Surgery 2012;12(10):227-231.
2. Knapp S. Solanum Section Geminata (Solanaceae). Floral Neotropica Monograph 2011;84(1):1-405.
3. Osei MK, Banful B, Osei CK, Oluoch MO. Characterization of African Eggplant for Morphological Characteristics., J. Agric. Sci. Technol. 2010;4(3):33-37.

4. Chinedu SN, Olasumbo AC, Eboji OK, Emiloju OC, Arinola OK, Dania DI. Proximate and Phytochemical Analyses of *Solanumaethiopicum Linn L.* and *Solanummacrocarpon L.* Fruits, Res.J.Chem.Sci.2011;1(3):24-35
5. Nwanna E, Ibukun E, Oboh G, Ademosun A, Athayde M. HPLC-DAD Analysis and in-vitro property of polyphenolss extracts from *Solanumaethiopicum Linn* fruits on  $\alpha$ -amylase,  $\alpha$ -glucosidase and angiotensin-1- converting enzyme activities. International journal of Biomedical sciences.2014;10(4):272-281
6. Anosike C, Obidoa O,Ezeanyika S. The anti-inflammatory activity of Garden egg on egg albumin-induced oedema and granuloma tissue formation in rats. Asian pacific Journal of Tropical Medicine:2012;5(1):62-66
7. Adedosu O, Badmus J, Adeleke G, Afolabi O, Adekunle A, Olabode F,Fakunle P. Oxidative damage and dietary antioxidants: The roles of extract and fractions of *Solanumaethioicum* leaves. Canadian Journal of Pure and Applied Sciences.2015;9(1):3185-3192
8. Emiloju EO,Chinedu SN. Effect of *Solanumaethiopicum Linn* and *Solanummacrocapon* Fruits on the Weight Gain, Blood Glucose and Liver Glycogen of Wister Rats. World Journal of Nutrition and Health.2016;4(1):1-4
9. Bello SO, Muhammad BY, Gammaniel KS, Abdu-Aguye I, Ahmed HH, Njoku HC et al. Preliminary Evaluation of the Toxicity and Some Pharmacological Properties of the Aqueous Crude Extract of *Solanummelongena*. Res J AgricBiol Sci. 2005;1(1):1–9.
10. Igwe SA, Akunyili DN, Ogbogu C. Effects of *Solanummelongena* (garden egg) on some visual functions of visually active Igbos of Nigeria. J Ethnopharmac. 2003;86(2–3):135–138 .
11. Sanchez–Mata MC, Yokoyama WE, Hong YJ, Prohens J. Alpha–solasonine and alpha–Solamargine Contents of Gboma (*Solanummacrocarpon L.*) and Scarlet (*Solanumaethiopicum Linn L.*) Eggplants. J Agric Food chem. 2010;58(9):5502–5508
12. Akinmoladun AC, Ibukun EO, Afor E, Obuotor EM, Farombi EO. Phytochemical constituent and antioxidant activity of extract from the leaves of *Ocimumgratissimum*. Sci. Res. Essay. 2007;2:63-166.

13. Zhou JR, Erdman JW Jr. Phytic acid in health and disease. *Critical Reviews in Food Science and Nutrition*. 1995;35(6):495–508
14. Nwanna E, Oboh G. Antioxidant and hepatoprotective properties of polyphenol extracts from *Telfairia occidentalis* (Fluted Pumpkin) leaves on acetaminophen induced liver damage. *Pak J Biol Sci*. 2007;10(16):2682-2687.
15. Shalom NC, Abayomi CO, Okwuchukwu KE, Opeyemi CE, Olajumoke KA, Damilola ID. Proximate and Phytochemical Analyses of *Solanumaethiopicum* Linn L. and *Solanummacrocarpon* L. Fruits. *Research Journal of Chemical Sciences*. 2011;1(3):63–70.
16. Asl MN, Hossein H. Review of pharmacological effects of *Glycyrrhiza* sp. and its bioactive compounds. *Phytotherapy Res*. 2008;22(6):709–724
17. Obidoa O, Joshua, Parker E, Eze NJ. Phytochemical Analysis of *Cocosnucifera* L. *Journal of Pharmacy Research*. 2010; 3(2):280-286.
18. Akinyemi OA, Oyelere FS. Phytochemical profile of selected morphological organs of *Cocosnucifera* I. *EJBPS* 2019; 6(11):54-58.
19. Sabri FZ, Belarbi M, SabriS, Alsayadi M. Phytochemical Screening and identification of some compounds from Mallow. *J. Nat. Prod. Plant Resour*. 2012;2(4):512-516.
20. Malliga E, Dhanarajan MS, Rajalakshmi A, Jayachitra A, Pardhasaradhi M, Narasimharao B. Analysis of Phytochemicals, Antibacterial and Antioxidant activities of *Moringaoleifera* Lam. Leaf extract- an in vitro study. *Int. J. Drug Dev. & Res*, 2014;6(4):173-180.
21. Ayoade ET, Akinyemi AO, Oyelere FS. Phytochemical profile of different morphological organs of *Moringaoleifera* plant. *J Phytopharmacol* 2019;8(6):295-298.
22. Eze SO, Kanu CQ. Phytochemical and nutritive composition analysis of *Solanumaethiopicum* L. *J Pharm Scilnnov*. 2014;3(4):358-362 <http://dx.doi.org/10.7897/2277-4572.034172>
23. Eletta OAA, Orimolade BO, Oluwaniyi OO, Dosumu OO. Evaluation of Proximate and Antioxidant Activities of Ethiopian Eggplant (*Solanumaethiopicum* LinnL) and Gboma Eggplant (*Solanummacrocarpon* L) *J. Appl. Sci. Environ. Manage*. 2017;21(5):967-972

24. Wink M, Schmeller T, Latz-Briining B. Modes of action of allelochemical alkaloids: Intraction with neuroreceptors, DNA and other molecular targets. *Journal of chemical Ecology*.1998;24:1888-1937.
25. Rao RVK, Ali N, Reddy MN. Occurrence of both saponins and alkaloid lycorine in *Curculigoorchioides*. *Indian Journal Pharma Science*. 1978;40:104-105.
26. Bassolé IHN, Juliani HR. Essential oils in combination and their antimicrobial properties. *Molecules*. 2012;17:3989–4006
27. Lang G, Buchbauer G. A review on recent research results (2008–2010) on essential oils as antimicrobials and antifungals. A review. *FlavourFragr. J.*2012;27:13–39.
28. Hajhashemi V, Ghannadi A, Pezeshkian SK. Antinociceptive and anti-inflammatory effects of *Saturejahortensis* L. extracts and essential oil. *J. Ethnopharmacol.* 2002;82:83–87.
29. Abdollahi M, Karimpour H, Monsef-Esfehani HR. Antinociceptive effects of *TeucriumPolium* L. total extract and essential oil in mouse writhing test. *Pharmacol. Res.* 2003;48:31–35.
30. Ross JA, Kassum CM. Dietary Saponins Availability, Metabolic Effects and Safety. *Journal of Annual Review on Nutrition* 2012; 24(3):19-30.
31. De Bruyne T, Pieters L, Deelstra H, Vlietinck A. Condensed vegetables tannins: biodiversity in structure and biological activities. *Biochemical System Ecology*. 1999;27:445–59.
32. Dolara P, Luceri C, De Filippo C, Femia AP, Giovannelli L, Carderni G, et al. Red wine polyphenols influence carcinogenesis, intestinal microflora, oxidative damage and gene expression profiles of colonic mucosa in F344 rats. *Mutation Research*, 2005;591:237–46.
33. Serrano J, Puupponen-Pimia R, Dauer A, Aura A, Saura-Calixto F. Tannins: current knowledge of food sources, intake, bioavailability and biological effects. *Molecular Nutrition Food Research*, 2009;53:S310–29.
34. McNaught KM. Phytochemical and Human Health. *Journal of the Italian Phamacological Society* 2008;55(3):207-216.