

**The prevalence of using prescribed &
unprescribed “Roaccutane (Isotretinoin)”
among females Riyadh, Saudi Arabia: A
cross sectional study**

**Dr. Mahmoud A Mahmoud¹, Reem A Almohaini², Raghad A Assiri²,
Reema A Alkhatabi^{2*}, Reema S Alobaishi², Rola F Alharbi²,**

*(Put * above the corresponding author and give telephone number, fax number and email ID in the footer)*

¹*Assistant professor, College of Medicine, Imam Mohammad ibn Saud Islamic University,
Riyadh, Saudi Arabia.*

²*College of Medicine, Imam Mohammad ibn Saud Islamic University, Riyadh, Saudi Arabia.*

ABSTRACT

Aims: To determine the prevalence of using prescribed and unprescribed Isotretinoin and to assess knowledge of its adverse effects among females in Riyadh, SA.

Study design: cross-sectional study.

Place and Duration of Study: online questionnaire collected throughout two months by distributing the questionnaire via social media platforms.

Methodology: We included 385 female patients (age range 15-45 years) use Roaccutane (Isotretinoin) with or without prescribed. Statistical analysis was performed using Rv 3.6.3. Counts and percentages was used to summarize the distribution of categorical variables and data. Chi-square test was used to estimate the association between categorical variables. Hypothesis testing was performed at a 5% level of significance.

Results: 568 respondents completed the questionnaire. Of these, 440 were from Riyadh. The majority of respondents were aged 15 - 20 (46.4%) and 21 - 25 (35.2%). Knowledge regarding some side effects of Roaccutane was significantly higher in respondents who reported using Roaccutane than those who did not. Knowledge of dry and cracked lips as a side effect was significantly higher in users (91.9%) than non-users (73.1%). Similarly, knowledge regarding dry eyes, depression, continuous thirst, nose dryness, bleeds, and effect of Roaccutane on liver biochemical profile, lipid profile, and bones was higher in users than non-users, regarding the teratogenic effect of Roaccutane was insignificantly different between groups.

Conclusion: Females in Riyadh are aware of the side effects of isotretinoin, particularly dryness of the skin, eyes, and lips. Yet there is insufficient knowledge regarding its teratogenic effect. Based on the reported results, a few participants obtained Roaccutane without a prescription from the doctor, and some obtained their information from unreliable sources (i.e., friends/relatives). Our results suggest good practice towards the prescription of isotretinoin.

Keywords: Isotretinoin, Roaccutane, Public health, Awareness, Side effects, Acne vulgaris

1. INTRODUCTION

Isotretinoin, also known as (Accutane or Roaccutane) is a synthetic vitamin A derivative, or retinoid, one of the most common and effective medications for treating acne vulgaris. Acne vulgaris (adolescent acne) is a chronic medical condition affecting the skin, precisely the pilosebaceous follicle, and is mainly characterized by alternating periods of exacerbation and stability, leading to polymorph cutaneous lesions that may leave scars after regression. [1-3] This common condition nearly affects all adolescents. The onset of the appearance of acne among females is usually between 18-20 years, or it may be delayed until 25-30 years. However, it can affect all ages as the risk increases with age, as mentioned in previous literature; in the pediatric population aged 10 to 12 years, 28% to 61% of the population has been clinically diagnosed with acne vulgaris. On the contrary, 79% to 95% of 16 to 18 years old adolescents are affected. [3-8] In 1982, Isotretinoin was proved by FDA (Food and Drug Administration) as a treatment for acne. [1][2] In addition, substantial studies recommended the use of Isotretinoin for treating acne more than other agents, as it is a highly effective agent. As known, this is not the only use of Isotretinoin; it can also act as a chemotherapeutic agent as it can treat various types of cancers, according to literature. [9] [10] Moreover, Previous studies showed that the use of Isotretinoin is associated with many side effects; the most observed and identified side effect was mucocutaneous such as retinoid dermatitis, severe retinoid cheilitis, epistaxis, dry eyes, or conjunctivitis, which are dose-dependent predictable side effects. Among adolescents, mood changes, including depression, are commonly reported; other psychiatric manifestations such as anxiety and suicidal thoughts are still controversial. [11-13] Teratogenicity is well known and regarded as one of the most severe and devastating adverse effects. According to FDA and previous literature, Isotretinoin causes congenital disabilities, abortion, and premature births. The recognized and known embryopathy pattern includes severe craniofacial, central nervous system, cardiovascular, and thymic malformations. Ear abnormalities include microtia, low set ears, and anotia; central nervous system defects include microcephalus, hydrocephalus, and reduction malformations of the brain; cardiovascular defects include transposition of the great arteries, tetralogy of Fallot, ventral septal defects, and aortic arch abnormalities. [11], [14-16] Due to this concern, the Saudi Food and Drug Authority (SFDA) started implementing a pregnancy prevention program (PPP) to raise the awareness of females regarding

the teratogenic effects of Isotretinoin. The program involves signing a medical consent form by the female user, contraceptive guide, and patient guide. The consent aims to prevent potential severe congenital disabilities that Isotretinoin may cause. The consent includes information and warnings about Isotretinoin use [17]. Through intensive literature, there was a couple of previous research with a similar aim around Saudi Arabia; However, there was no study conducted in the capital of Saudi Arabia that Measure the awareness of females and their knowledge about side effects of Isotretinoin, also the prevalence of using prescribed and unprescribed of this type of treatment.

2. METHODS

This study will utilize a cross-sectional design using an electronic **validated Arabic questionnaire** distributed over multiple social media platforms. The sample size is estimated to be 385 participants, large enough to reach a confidence interval of 95% as calculated using the Raosoft Sample Size Calculator (Raosoft, Inc. Seattle, WA). **Ethical approval for this study was obtained from the Ethics Committee in Imam Mohammad ibn Saud Islamic University.** The data will be collected throughout two months by distributing the survey over social media platforms. The instrument is derived from an intensive literature review. **After developing the questionnaire, it will be translated into Arabic and validated via running a pilot testing on twenty random participants to recognize any issues related to context and content.** The questionnaire comprises 16 questions (yes/no questions, multiple choices, short answers, and check-boxes). Those questions were divided into two sections: section A was related to demographic data regarding the participant; section B was to assess their knowledge regarding the use of Isotretinoin and the method of prescription. **Also, we asked them about the lab investigations they did before the usage of Roaccutane.** All participants will receive written consent before completing the questionnaire.

2.1 Statistical analysis

Statistical analysis will be performed using R v 3.6.3. Counts and percentages will be used to summarize the distribution of categorical variables. A Chi-square test will be used to estimate the association between categorical variables. Hypothesis testing will be performed at a 5% level of significance. Data will be summarized using counts and percentages.

2.2 Inclusion criteria

1. Any Female participants willing to fill the survey.
2. Living in Riyadh, Saudi Arabia.
3. Age 15 years and above.

2.3 Exclusion criteria

1. Participants younger than 15 years,
2. Currently living outside of Riyadh Saudi Arabia
3. Participants with disabilities that prevents them from filling the survey

3. RESULTS AND DISCUSSION

Table 1. Sociodemographic characteristics of the included respondents

	[ALL]	N
	<i>N=440</i>	
Age:		440
15-20	204	
21-25	155	
26-30	29 (6.59%)	
31-35	22 (5.00%)	
41-45	16 (3.64%)	
45+	14 (3.18%)	
Marital status:		440
Divorced	7 (1.59%)	
Married	63 (14.3%)	
Single	369	
Widowed	1 (0.23%)	
Highest educational		440
Middle school	4 (0.91%)	
High school	125	
Bachelor	293	
Post-graduate	18 (4.09%)	
Data were summarized using counts and		

Five hundred sixty-eight respondents completed the questionnaire. Of these, 440 were from Riyadh. The majority of respondents were aged 15 - 20 (46.4%) and 21 - 25 (35.2%). Single respondents represented 83.9% of the study sample, and 14.3% were married. Regarding education, 66.6% of the respondents had a bachelor's degree, and 28.4% completed only high school.

Table 2. prevalence and awareness regarding the use of Roaccutane

	[ALL]	N
	<i>N=440</i>	
Ever used Roaccutane		440
No	279	
Yes	161	
Any relatives use Roaccutane		161
No	23 (14.3%)	
Yes	138	
Source of information regarding Roaccutane and its		161
Dermatologist	79 (49.1%)	
Friends and relatives	74 (46.0%)	
Social media	8 (4.97%)	
Roaccutane prescribed by a physician		161
No (Available without prescription)	3 (1.86%)	
Yes	158	
Laboratory investigations performed before using		161
No	8 (4.97%)	
Yes	153	
Duration of Roaccutane use:		161
< 2 months	10 (6.21%)	
2 - 3 months	21 (13.0%)	
3 - 5 months	65 (40.4%)	
6 - 12 months	49 (30.4%)	
1+ year	16 (9.94%)	
Roaccutane use:		161
Continuous	137	
Intermittent	24 (14.9%)	
Aware of Roccutane's side effects:		161
No	10 (6.21%)	
Yes	151	
Data were summarized using counts and percentages		

One-third of the respondents reported using Roaccutane (n = 161, 36.6%) and 63.4% did not. Of those who used Roaccutane, 85.7% reported knowing relatives who used it. The source of information regarding Roaccutane and its uses varied between respondents. The dermatologist was the main source of information in

49.1% of the cases, while friends and relatives were the main sources in 46%. Social media was the main source of information in the remaining 4.97% of the respondents. All but three respondents (98.1%) mentioned that a physician prescribed Roaccutane. The great majority of the respondents reported doing laboratory investigations before using Roaccutane (95%). The duration of use varied between respondents, with only 6.21% and 9.94% reporting its use for < 2 months and 1+ year, respectively. The majority of the respondents reported using Roaccutane for 3 - 5 months (40.4%) and 6 - 12 months (30.4%), respectively. The use of Roaccutane was continuous in 85.1% of the cases and intermittent in the remaining 14.9%. Most of the respondents were aware of the side effects of Roaccutane (93.8%).




Figure 1. Awareness regarding side effects of Roaccutane

All respondents were asked to identify the side effects of Roaccutane (n = 440). The most commonly identified side effects of Roaccutane were dry skin (84.5%), dry/cracked lips (80%), and dry eyes (33.6%). The least commonly identified side effects were diplopia (5.7%), hypohidrosis (13.4%), and nausea/loss of appetite (13.9%).




Figure 2. Knowledge regarding the teratogenic effect of isotretinoin stratified by marital status

Results showed that knowledge regarding the teratogenic effect of Isotretinoin was low across all age groups and was lowest in respondents aged 15 - 20 years old (n = 440).




Figure 3. Laboratory investigations performed before using Roaccutane

Respondents who used Roaccutane were asked to identify the laboratory investigations performed before its use (n = 161). Liver biochemical profile was performed by 78.9% of the users. Triglycerides and cholesterol levels were checked by 69.6% of the respondents, and CBC was performed by 64.6% of the users. Pregnancy testing was performed by 3.7% of the users.

Table 3. Association between Roaccutane use and its side effects

	Non-users	Users	p. overall
	N=279	N=161	
Dry skin:	231 (82.8%)	141 (87.6%)	0.230
Dry and cracked lips:	204 (73.1%)	148 (91.9%)	<0.001
Dry eyes:	162 (58.1%)	136 (84.5%)	<0.001
Depression:	78 (28.0%)	73 (45.3%)	<0.001
Eczema:	34 (12.2%)	28 (17.4%)	0.171
Continuous thirst:	144 (51.6%)	65 (40.4%)	0.030
Nose dryness and nosebleeds:	84 (30.1%)	97 (60.2%)	<0.001
Elevated liver enzymes:	67 (24.0%)	66 (41.0%)	<0.001
Hyperlipidemia and hypertriglyceridemia:	45 (16.1%)	70 (43.5%)	<0.001
Myalgia and arthralgia:	66 (23.7%)	88 (54.7%)	<0.001
Insomnia:	31 (11.1%)	19 (11.8%)	0.949
Hair loss:	98 (35.1%)	79 (49.1%)	0.006
Skin eruption (Face):	36 (12.9%)	13 (8.07%)	0.163
Fatigue and headache:	55 (19.7%)	64 (39.8%)	<0.001
Teratogenicity (pregnancy):	154 (55.2%)	96 (59.6%)	0.422
Nausea and loss of appetite:	38 (13.6%)	23 (14.3%)	0.959
Hypohidrosis:	43 (15.4%)	16 (9.94%)	0.139
Diplopia:	16 (5.73%)	9 (5.59%)	1.000

**Counts and percentages were used to summarize the responses*

**statistical analysis was performed using Chi-square test of independence*

Results showed that knowledge regarding some side effects of Roaccutane was significantly higher in respondents who reported using Roaccutane than those who did not. Knowledge regarding dry and cracked lips was significantly higher in users (91.9%) than non-users (73.1%). Similarly, knowledge regarding dry eyes, depression, continuous thirst, nose dryness, and bleeds was higher in users than non-users. Knowledge regarding the effect of Roaccutane on liver biochemical profile, lipid profile, and bones was also higher in users than non-users. Knowledge regarding the teratogenic effect of Roaccutane was not significantly different between groups.

Emotional, Physical, and scarring are significant burdens of acne in patients of various ages. Effective regimens are readily available. However, consistent and accurate use of these regimens is required for effective disease management, reduced risk for scarring, and enhancing various factors that affect the quality of life [18]. The rise in hormonal levels during puberty causes the formation of acne. The presence of other diseases such as polycystic ovary syndrome (PCOS) can also affect the appearance of acne [19]. Sensitive skin, lifestyle-related stress, using full-coverage foundations, and discontinuation of oral contraceptives increase the severity of acne [20]. Roaccutane or Isotretinoin is an oral vitamin A derivative, considered one of the most effective options for managing severe acnes [21]. Advantages of Isotretinoin include cost-effectiveness, especially when compared with other medications used to treat severe acne. However, it is teratogenic when used during pregnancy and can cause congenital disabilities, such as severe

cardiovascular, craniofacial, thymic, and central nervous system malformations [16]. Patients who suffer from acne know about isotretinoin drugs in one way or other. However, some patients may hesitate from starting this drug due to a lack of information about the drug. Proper understanding and awareness regarding the drug and its side effects help minimize adverse effects and increase compliance. The current study aimed to assess the use and awareness regarding the side effects of Isotretinoin in female patients in Riyadh. A similar study was conducted in Riyadh recently. However, the distribution of age was similar to that observed in the current study. In the study conducted by Ibrahim and colleagues, the majority of the respondents were aged 20 - 30 years old [22]. In this study, one-half of the respondents were aged 15 - 20 years old, and 42% were aged 21 - 30 years old, which adds to the existing knowledge regarding Isotretinoin and awareness regarding its side effects. Nonetheless, a discrepancy was observed in the knowledge regarding the teratogenic effect of Isotretinoin. A previous study was done in Riyadh showed that ~90% of the respondents were aware of the side effects of Isotretinoin [22]. Also, a study conducted in Eastern Saudi Arabia showed that 88% of the respondents were aware of the teratogenic effect of Isotretinoin [23]. The results in this study were not comparable to these two previous studies, which showed that 56.8% of the respondents were not aware of the teratogenic effect of Isotretinoin [22]. The insufficient knowledge regarding the teratogenic effect of Roaccutane can be explained by the high number of single participants included in this study (83%) compared to other studies and the low number of isotretinoin users included in the current study (36%). Thus, it can be expected that knowledge regarding Isotretinoin and its teratogenic effect would be higher if the users were married, of childbearing age, previously used, or are currently using Isotretinoin as was the case in a cross-sectional study conducted in Makkah [24]. Another explanation would be the source of information regarding Isotretinoin, with 50% of the respondents relying on relatives/friends or social media as a source of information. Nonetheless, our results are also consistent with those reported in another cross-sectional study in Western Saudi Arabia which showed even lower knowledge regarding the teratogenic effect of Isotretinoin [25]. The similarity in knowledge can be explained by the similarities in the demographic characteristics

(marital status and age), particularly the lower age of the included participants. These results suggest that knowledge regarding the teratogenic effect of Isotretinoin is still suboptimal, a finding which is supported by evidence from another recent cross-sectional study [26]. Dryness of the skin, eyes, and lips were the top three side effects identified by the respondents, followed by nose dryness and bleeding. These results are similar to those reported in several cross-sectional studies [24,25,27], where the dryness of the eyes, lips, and skin were the most common side effects. On the other hand, the least experienced side effect was depression; this result is consistent with what was observed in previous studies [13,28]. In this study, most respondents who used Isotretinoin (95%) performed at least one of the needed laboratory investigations before using Isotretinoin. A study conducted in 2019 in Eastern Saudi Arabia showed that half of the participants (44.2%) used Isotretinoin with only mild acne as their first choice, which is not consistent with the current guidelines. The study also showed that one-fifth of the respondents did not assess the blood glucose level, lipid profile, liver enzymes before starting isotretinoin [23]. These results are in line with what was observed in this current study. One-third of the respondents did not assess cholesterol or triglyceride levels, one-third did not perform a complete blood count (CBC), and one-fifth did not perform a liver biochemical profile. Interestingly, only 3.7% of the respondents did pregnancy testing before using the medication. A study conducted in Riyadh found that knowledge regarding the teratogenic effects of Isotretinoin was low and recommended improvement in the risk reduction counseling skills [29]. Many women using Isotretinoin do not monitor for pregnancy before and during the use of Isotretinoin. Only 44%-67% of women reported doing a pregnancy test before starting Isotretinoin, and only 20% reported taking a repeated pregnancy test [30]. Moreover, most women use over-the-counter (OTC) pregnancy tests for human chorionic gonadotropin (HCG) hormone, which is characterized by low sensitivity and may cause false-negative results, especially if done shortly after conception [31]. Repeat testing may help detect pregnancy and earlier termination of treatment or pregnancy [32]. We suggest that the difference regarding the teratogenic effect of Isotretinoin can somewhat be explained by the source of information regarding Isotretinoin, where dermatologists were the primary source of information in only

50% of the included respondents. At the same time, friends, relatives, and social media were the primary source in the remaining 50%. It has also been suggested that a contraceptive information sheet can significantly improve the knowledge of the patients regarding contraception and may prevent isotretinoin-induced congenital disabilities [33]. The knowledge regarding the side effects of Isotretinoin was significantly higher in users than non-users previously reported in the literature [12,25]. With so many patients taking medicines for a variety of ailments, it's critical for doctors to know what prescriptions their patients are taking, why they're taking them, and if they're suffering any adverse effects.

4. CONCLUSION

Results suggest that women in Riyadh, Saudi Arabia, are somewhat aware of the side effects of isotretinoin, particularly dryness of the skin, eyes, and lips. Results also showed insufficient knowledge regarding its teratogenic effect. Based on the reported results, some of the included patients did not do the needed tests such as pregnancy tests and blood tests before starting isotretinoin. Moreover, some study participants obtained Roaccutane without a prescription from the doctor, and some obtained their information from unreliable sources (i.e., friends/relatives ,internet). In conclusion, our results suggest good practice towards the prescription of isotretinoin. Knowledge regarding isotretinoin and its use should be disseminated in the community to prevent possible teratogenicity from its use. Therefore, we encourage rising awareness regarding Saudi FDA pregnancy prevention program to prevent future incidents.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

Authors may use the following wordings for this section: "Author A' designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. 'Author B' and 'Author C' managed the analyses of the study. 'Author C' managed the literature searches..... All authors read and approved the final manuscript."

REFERENCES

- 1.Ramos-e-Silva M, Carneiro SC. Acne vulgaris: review and guidelines. *Dermatol Nurs.* 2009;21(2):63-69.
- 2.Hauk L. Acne Vulgaris: Treatment Guidelines from the AAD. *Am Fam Physician.* 2017;95(11):740-741.
- 3.Aydemir EH. Acne vulgaris. *Turk Pediatri Ars.* 2014;49(1):13-16.
- 4.Huang YC, Cheng YC. Isotretinoin treatment for acne and risk of depression: A systematic review and meta-analysis. *J Am Acad Dermatol.* 2017;76(6)
- 5.Kutlubay Z, Kecici AS, Engin B, Serdaroglu S, Kutlubay Z, Kecici AS. Acne Vulgaris Acne Vulgaris. 2017;11:247–55.
- 6.Rademaker M, Garioch JJ, Simpson NB. Acne in schoolchildren: no longer a concern for dermatologists. *BMJ.* 1989;298(6682):1217-1219
- 7.Merlin K, Kilkenny M, Plunkett A, Marks R. The prevalence of common skin conditions in Australian school students: 4 Tinea pedis. *Br J Dermatol.* 1999;140(5):897–901.
- 8.Lello J, Pearl A, Arroll B, Yallop J, Birchall NM. Prevalence of acne vulgaris in Auckland senior high school students. *N Z Med J.* 1995;108(1004):287–9.
- 9.Alkhalidy HY, Assiri AM, Fatima S, Owaidah T. Isotretinoin is active in the initial management of acute pro-myelocytic leukemia. *Leuk Res Rep.* 2020;14(100220):100220.
- 10.Khuri FR, Lee JJ, Lippman SM, Kim ES, Cooper JS, Benner SE, et al. Randomized phase III trial of low-dose isotretinoin for prevention of second primary tumors in stage I and II head and neck cancer patients. *J Natl Cancer Inst.* 2006;98(7):441–50.
- 11.Layton A. The use of isotretinoin in acne. *Dermatoendocrinol.* 2009;1(3):162–9.
- 12.Molla A, Alrizqi HA, Alruhaili EMS, Alrizqi SA, Alsubhi AM. Assessment of knowledge, attitude, and practice in relation to use of isotretinoin among Al-Madinah population, Saudi Arabia. *IJMDC.* (2020), [cited October 02, 2021]; 4(1): 107-112

13. Ahmad HM. Analysis of clinical efficacy, side effects, and laboratory changes among patients with acne vulgaris receiving single versus twice daily dose of oral isotretinoin: Oral isotretinoin for acne vulgaris. *Dermatol Ther.* 2015;28(3):151–7.
14. Pastuszek A, Koren G, Rieder MJ. Use of the Retinoid Pregnancy Prevention Program in Canada: patterns of contraception use in women treated with isotretinoin and etretinate. *Reprod Toxicol.* 1994;8(1):63–8.
15. Mondal D, R Shenoy S, Mishra S. Retinoic acid embryopathy. *Int J Appl Basic Med Res.* 2017;7(4):264–5
16. Malvasi A, Tinelli A, Buia A, De Luca GF. Possible long-term teratogenic effect of isotretinoin in pregnancy. *Eur Rev Med Pharmacol Sci.* 2009;13(5):393–6.
17. Saudi Food and Drug Authority, 2020b. Isotretinoin Medical Consent Form. https://www.sfd.gov.sa/en/drug/about/sector_departments/national_pharmacovigilance_center/Documents/isotretinoin-second-version-of-RMM-%283%29.pdf
18. Zaenglein AL Md. Psychosocial issues in acne management: Disease burden, treatment adherence, and patient support. *Semin Cutan Med Surg.* 2015;34(5 Suppl):S92-4.
19. Skroza N, Tolino E, Proietti I, Bernardini N, La Viola G, Nicolucci F, et al. Women and acne: any difference from males? A review of the literature. *G Ital Dermatol Venereol.* 2016;151(1):87–92.
20. Chlebus E, Chlebus M. Factors affecting the course and severity of adult acne. Observational cohort study. *J Dermatolog Treat.* 2017;28(8):737–44.
21. Layton AM, Dreno B, Gollnick HPM, Zouboulis CC. A review of the European Directive for prescribing systemic isotretinoin for acne vulgaris: Editorial. *J Eur Acad Dermatol Venereol.* 2006;20(7):773–6.
22. Ibrahim AAM, Alshatri AA, Alsuwaidan S, Almutairi L, Aljasser N, Mahmoud MA, et al. Awareness of isotretinoin use and Saudi FDA pregnancy prevention program in Riyadh, Saudi Arabia: A cross-sectional study among female patients. *Saudi Pharm J.* 2021;29(6):527–32.
23. Younis NS, Al-Harbi NY. Public understanding and awareness of isotretinoin use and safety in Al ahsa, eastern Saudi Arabia. *Ther Innov Regul Sci.* 2019;53(5):618–22.
24. Al-Mekhlafi R, Attiyah RE, Haddad YR, Salah LA. Awareness on teratogenic effects of isotretinoin and compliance with precautionary measures among women of childbearing age in Makkah Province, Saudi Arabia. *Obstet Gynecol Int.* 2021;2021:9966300.
25. Bakheet KMA, Alghanemi RG, Alsiyoufi AM, Abduljabbar M, Hariri J. Females' knowledge and use of isotretinoin (roaccutane) in the western region of Saudi Arabia. *Cureus.* 2020;12(12):e12148.
26. Alharbi M. Challenges and barriers to start isotretinoin in acne patients. *Australasian Medical Journal [Internet].* 2020;13(4).
27. Al-Harbi M. Concerns and awareness of acne patients about isotretinoin in qassim region of saudi arabia. *Int J Health Sci (Qassim).* 2010;4(1):47–51.

28. Agarwal US, Besarwal RK, Bhole K. Oral isotretinoin in different dose regimens for acne vulgaris: a randomized comparative trial. *Indian J Dermatol Venereol Leprol*. 2011;77(6):688–94.
29. Alharithy R, Algoblan S, Bakhsh S. Women's experiences regarding isotretinoin risk reduction counseling in Riyadh. *J dermatol dermatol surg*. 2019;23(1):13.
30. Robertson J, Polifka JE, Avner M, Chambers C, Delevan G, Koren G, et al. A survey of pregnant women using isotretinoin. *Birth Defects Res A Clin Mol Teratol*. 2005;73(11):881–7.
31. Cole LA, Ladner DG. Background hCG in non-pregnant individuals: need for more sensitive point-of-care and over-the-counter pregnancy tests. *Clin Biochem*. 2009;42(3):168–75.
32. Abrams L, Maibach E, Lyon-Daniel K, Feldman SR. What is the best approach to reducing birth defects associated with isotretinoin? *PLoS Med*. 2006;3(11):e483.
33. Leger MC. Promoting safe use of isotretinoin by increasing contraceptive knowledge. *JAMA Dermatol*. 2015;151(4):393–4.

APPENDIX

